

## THE NONSPLIT DOMINATION NUMBER OF A GRAPH

V. R. KULLI AND B. JANAKIRAM

*Department of Mathematics, Gulbarga University, Gulbarga 585 106, India*

*(Received 3 April 1996; after final revision 9 February 1998; Accepted 7 December 1999)*

A dominating set  $D$  of graph  $G = (V, E)$  is a nonsplit dominating set if the induced subgraph  $\langle V - D \rangle$  is connected. The nonsplit domination number  $\gamma_{ns}(G)$  of  $G$  is the minimum cardinality of a nonsplit dominating set. In this paper, many bounds on  $\gamma_{ns}(G)$  are obtained and its exact values for some standard graphs are found. Also its relationship with other parameters is investigated.

**Key Words :** Nonsplit Domination Number; Graph

### 1. INTRODUCTION

The graphs considered here are finite, undirected nontrivial and connected without loops or multiple edges.

Let  $G = (V, E)$  be a graph. A set  $D \subseteq V$  is a dominating set of  $G$  if every vertex in  $V - D$  is adjacent to some vertex in  $D$ . The domination number  $\gamma(G)$  of  $G$  is the minimum cardinality of a dominating set.

A dominating set  $D$  of  $G$  is a connected dominating set if the induced subgraph  $\langle D \rangle$  is connected. The connected domination number  $\gamma_c(G)$  of  $G$  is the minimum cardinality of a connected dominating set.

Recently Kulli and Janakiram introduced the concept of Split Domination<sup>5</sup>.

A dominating set  $D$  of a graph  $G = (V, E)$  is a split dominating set if the induced subgraph  $\langle V - D \rangle$  is disconnected. The split domination number  $\gamma_s(G)$  of  $G$  is the minimum cardinality of a split dominating set.

The reader is referred to [1], [2], [3] for survey of results on domination.

Any undefined term in this paper may be found in Harary<sup>4</sup>. Unless stated, the graph has  $p$  vertices and  $q$  edges.

The purpose of this paper is to introduce the concept of Nonsplit Domination.

A dominating set  $D$  of a graph  $G = (V, E)$  is a nonsplit dominating set if the induced subgraph  $\langle V - D \rangle$  is connected. The nonsplit domination number  $\gamma_{ns}(G)$  of  $G$  is the minimum cardinality of a nonsplit dominating set.

We call a set of vertices a  $\gamma$ -set if it is a dominating set with cardinality  $\gamma(G)$ . Similarly, a  $\gamma_c$ -set, a  $\gamma_s$ -set and a  $\gamma_{ns}$ -set are defined.

## 2. RESULTS

We start with some elementary results. Since their proofs are trivial, we omit the same.

**Theorem 1** — For any graph  $G$ ,

$$\chi(G) \leq \gamma_{ns}(G). \quad \dots (1)$$

**Theorem 2** — For any graph  $G$ ,

$$\chi(G) = \min \{ \gamma_s(G), \gamma_{ns}(G) \}. \quad \dots (2)$$

In<sup>3</sup>, Cockayne and Hedetniemi gave necessary and sufficient conditions for a minimal dominating set.

**Theorem A<sup>3</sup>** — A dominating set  $D$  of a graph  $G$  is minimal if and only if for each vertex  $v \in D$  one of the following conditions is satisfied.

(i) there exists a vertex  $u \in V - D$  such that  $N(u) \cap D = \{v\}$ ;

(ii)  $v$  is an isolated vertex in  $\langle D \rangle$ .

**Theorem 3** — A nonsplit dominating set  $D$  of  $G$  is minimal if and only if for each vertex  $v \in D$  one of the following conditions is satisfied :

(i) there exists a vertex  $u \in V - D$  such that  $N(u) \cap D = \{v\}$ ;

(ii)  $v$  is an isolated vertex in  $\langle D \rangle$ ;

(iii)  $N(v) \cap (V - D) = \phi$ .

PROOF : Suppose  $D$  is minimal. On the contrary, if there exists a vertex  $v \in D$  such that  $v$  does not satisfy any of the given conditions, then by Theorem A,  $D' = D - \{v\}$  is a dominating set of  $G$  by (iii),  $\langle V - D' \rangle$  is connected. This implies that  $D'$  is a nonsplit dominating set of  $G$ , a contradiction. This proves the necessity.

Sufficiency is straightforward.

Next we obtain a relationship between  $\gamma_{ns}(G)$  and  $\gamma_{ns}(H)$  where  $H$  is any spanning subgraph of  $G$ . We omit the proof.

**Theorem 4** — For any spanning subgraph  $H$  of  $G$ ,

$$\gamma_{ns}(G) \leq \gamma_{ns}(H). \quad \dots (3)$$

In the following two results, we obtain lower and upper bounds on  $\gamma_{ns}(G)$  respectively.

**Theorem 5** — For any graph  $G$ ,

$$\gamma_{ns}(G) \geq (2p - q - 1)/2. \quad \dots (4)$$

PROOF : Let  $D$  be a  $\gamma_{ns}$  - set of  $G$ . Since  $\langle V - D \rangle$  is connected,

$$q \geq |V - D| + |V - D| - 1.$$

This proves (4).

**Theorem 6** — For any graph  $G$ ,

$$\gamma_{ns}(G) \leq p - \omega(G) + 1 \quad \dots (5)$$

where  $\omega(G)$  is the clique number of  $G$ .

PROOF : Let  $S$  be a set of vertices of  $G$  such that  $\langle S \rangle$  is complete with  $|S| = \omega(G)$ . Then for any  $u \in S$ ,  $(V - S) \cup \{u\}$  is a nonsplit dominating set of  $G$ .

Thus (5) holds.

Now we list the exact values of  $\gamma_{ns}(G)$  for some standard graphs.

*Proposition 7* —

(i) For any complete graph  $K_p$  with  $p \geq 2$  vertices,

$$\gamma_{ns}(K_p) = 1. \quad \dots (6)$$

(ii) For any complete bipartite graph  $K_{m,n}$  with  $2 \leq m \leq n$ ,

$$\gamma_{ns}(K_{m,n}) = 2. \quad \dots (7)$$

(iii) For any cycle

$$C_p, \gamma_{ns}(C_p) = p - 2. \quad \dots (8)$$

(iv) For any wheel

$$W_p, \gamma_{ns}(W_p) = 1. \quad \dots (9)$$

(v) For any path  $P_p$  with  $p \geq 3$  vertices,

$$\gamma_{ns}(P_p) = p - 2. \quad \dots (10)$$

Our next result sharpens the inequality (5) for trees.

*Theorem 8* — If  $T$  is a tree which is not a star, then,

$$\gamma_{ns}(T) \leq p - 2. \quad \dots (11)$$

PROOF : Since  $T$  is not a star, there exist two adjacent cut vertices  $u$  and  $v$  with  $\deg u, \deg v \geq 2$ . This implies that  $V - \{u, v\}$  is a nonsplit dominating set of  $T$ .

Thus (11) holds.

*Theorem 9* — If  $\kappa(G) > \beta_0(G)$ , then

$$\gamma_{ns}(G) = \chi(G) \quad \dots (12)$$

where  $\kappa(G)$  is the connectivity of  $G$  and  $\beta_0(G)$  is the independence number of  $G$ .

PROOF : Let  $D$  be a  $\gamma$ -set of  $G$ . Since  $\kappa(G) > \beta_0(G) \geq \chi(G)$ , it implies that  $\langle V - D \rangle$  is connected. This proves that  $D$  is a  $\gamma_{ns}$ -set of  $G$ . Hence (12) follows.

**Theorem 10** — Let  $D$  be a  $\gamma_{ns}$  - set of a connected graph  $G$ . If no two vertices in  $V - D$  are adjacent to a common vertex in  $D$ , then

$$\gamma_{ns}(G) + \varepsilon(T) \geq p \quad \dots (13)$$

where  $\varepsilon(T)$  is the maximum number of endvertices in any spanning tree  $T$  of  $G$ .

PROOF : Let  $D$  be a  $\gamma_{ns}$  - set of  $G$ , given in the hypothesis. Since for any two vertices  $u, v \in V - D$ , there exist two vertices  $u_1, v_1 \in D$  such that  $u_1$  is adjacent to  $u$  but not to  $v$  and  $v_1$  is adjacent to  $v$  but not to  $u_1$ , this implies that there exists a spanning tree  $T$  of  $\langle V - D \rangle$  in which each vertex of  $V - D$  is adjacent to a vertex of  $D$ . This proves that  $\varepsilon(T) \geq |V - D|$ .

Thus (13) holds.

**Theorem 11** — If  $\delta(G) + \omega(G) \geq p + 1$ , then

$$\gamma_c(G) + \gamma_{ns}(G) \leq p \quad \dots (14)$$

where  $\delta(G)$  is the minimum degree of  $G$ .

PROOF : By (5),  $\gamma_{ns}(G) \leq p - \omega(G) + 1 \leq \delta(G)$ .

Let  $D$  be a  $\gamma_{ns}$  - set of  $G$ . Then every vertex in  $D$  is adjacent to some vertex in  $V - D$ . Thus  $\langle V - D \rangle$  is a connected dominating set of  $G$ , since  $\langle V - D \rangle$  is connected. This proves (14).

In the next result we obtain another upper bound on  $\gamma_{ns}(G)$ .

**Theorem 12** — For any graph  $G$ ,

$$\gamma_{ns}(G) \leq p - \text{diam}(G) + h + 1 \quad \dots (15)$$

where  $\text{diam}(G)$  is the diameter of  $G$  and  $h$  is the minimum number of vertices in a  $\gamma_{ns}$ -set of  $G$  which lie in between shortest  $u$ - $v$  path and  $d(u, v) = \text{diam}(G)$ .

PROOF : Let  $\text{diam}(G) = k$ . We consider the following cases.

Case 1 — Suppose  $u, v \in V - D$ . Then  $V - D$  has at least  $k + 1$  vertices.

Case 2 — Suppose  $u \in D$  and  $v \in V - D$ . If there exists a vertex  $u_1 \in V - D$  such that  $u_1$  is connected to  $u$  through the vertices of  $D$  then,  $d(u_1, v) \geq k - (h + 1)$  and hence  $V - D$  has at least  $k - h$  vertices. For otherwise, for every vertex  $u_1 \in V - D$  there exists a vertex  $w$  adjacent to  $u_1$  such that  $d(u, w) = d(u, v) + d(v, u_1) + d(u_1, w) \geq k + 1$ , a contradiction.

This implies that  $V - D = \{v\}$  and hence  $G = K_2$  or  $K_{1,2}$ .

Case 3 — Suppose  $u, v \in D$ . If there exist two vertices  $u_1, v_1 \in V - D$  such that  $u$  is connected to  $u_1$  and  $v$  is connected to  $v_1$  through the vertices of  $D$ , then  $d(u_1, v_1) \geq k - (h + 2)$  and hence  $V - D$  has at least  $k - h - 1$  vertices. For otherwise, there exists exactly one vertex  $u_1 \in V - D$  which is adjacent to both  $u$  and  $v$  and  $\{u_1\} = (V - D)$ . This implies that  $G$  is a star with at least three vertices.

Thus from the above all the three cases, it follows that  $V - D$  has at least  $k - h - 1$  vertices and hence (15) follows.

Now we obtain a lower bound on  $\gamma_{ns}(T)$ .

**Theorem 13** — For any tree  $T$ ,

$$\gamma_{ns}(T) \geq p - m \quad \dots (16)$$

where  $m$  is the number of vertices adjacent to endvertices.

PROOF : If  $T$  is  $K_2$ , the result is trivial. If  $T$  has at least three vertices and  $D$  is a  $\gamma_{ns}$  - set of  $T$ , then each vertex of  $V - D$  is a cutvertex of  $T$ . Let  $S$  be the set of all cutvertices which are adjacent to endvertices with  $|S|=m$ . Let  $u \in V - D$ . If  $u \in S$ , then  $D = V - S$  and (16) holds. If  $u \notin S$ , then there exists a cutvertex  $v \in D$  adjacent to  $u$ . Further, all vertices which are connected to  $v$  not through  $u$  also belonging to  $D$ . This implies that  $V - D$  has at most  $m$  vertices and (16) holds.

**Corollary 13.1** — For any tree  $T$ ,

$$\gamma_c(T) \leq \gamma_{ns}(T). \quad \dots (17)$$

Further if  $T$  is a path, then equality holds.

PROOF : If  $T$  has no cut vertices, then  $T=K_2$  and hence  $\gamma_c(T) = \gamma_{ns}(T) = 1$ .

Let  $S$  be the set of all cut vertices of  $T$  with  $|S|=p_1$  and  $S_1 \subseteq S$  be the set of all cut vertices such that each vertex of  $S_1$  is adjacent to an endvertex with  $|S_1|=p_2$ .

Thus,  $V(T) = p \geq p_1 + p_2$ .

Due to Sampath Kumar and Walikar<sup>6</sup>,

$$\gamma_c(T) = p_1.$$

Hence (17) follows from (16).

If  $T$  is path with  $p \geq 3$  vertices, then by (10) and the fact that  $\gamma_c(T) = p_1$ , the equality holds.

Next we obtain an upper bound on  $\gamma_{ns}(T)$ .

**Theorem 14** — For any tree  $T$ ,

$$\gamma_{ns}(T) \leq p - \max_v \{ \deg v - |e(v)| \}$$

where  $e(v)$  is the set of all endvertices adjacent to  $v$ .

PROOF : Let  $v$  be a vertex with  $\deg v - |e(v)|$  being maximum. Let  $u \in N(v)$ . Then it follows that  $V - N[v] \cup e(v) \cup \{u\}$  is a nonsplit dominating set of  $T$ . Hence (18) holds.

**Corollary 14.1** — For any tree  $T$ ,

$$\gamma_{ns}(T) \leq p - \Delta(T) + p_0 \quad \dots (19)$$

where  $\Delta(T)$  is the maximum degree of  $T$  and  $p_0$  is the maximum number of endvertices adjacent to a vertex of maximum degree.

**Corollary 14.2** — For any graph  $G$ ,

$$\gamma_{ns}(G) \leq p - \max_v \{ \deg v - |e(v)| \} \quad \dots (20)$$

where  $e(v)$  is the set of all vertices which are adjacent to  $v$  but not adjacent to any vertex of  $V - N(v)$ .

PROOF : This follows from the fact that for any  $v \in V$ , there exists a spanning tree  $T$  such that  $\deg_G v = \deg_T v$  and from (3) and (18).

The next result relates to  $\gamma_{ns}(G)$  and  $\gamma_s(\overline{G})$  where  $\overline{G}$  is the complement of  $G$ .

**Theorem 15** — *If  $\text{diam}(G) = 5$ , then*

$$\gamma_s(G) \geq \gamma_{ns}(\overline{G}). \quad \dots (21)$$

PROOF : Let  $D$  be a  $\gamma_s$ -set of  $G$ . Then every vertex in  $V - D$  is not adjacent to at least one vertex in  $D$ , since  $\text{diam}(G) = 5$ . Thus  $D$  is a dominating set of  $\overline{G}$  and further it is a nonsplit dominating set of  $\overline{G}$ , as  $\langle V - D \rangle$  is connected in  $\overline{G}$ .

This proves (21).

The following result is obvious. Hence we omit its proof.

**Theorem 16** — *Let  $G$  be a graph such that both  $G$  and  $\overline{G}$  are connected. Then*

$$(i) \quad \gamma_{ns}(G) + \gamma_{ns}(\overline{G}) \leq 2(p - 2); \quad \dots (22)$$

$$(ii) \quad \gamma_{ns}(G) \cdot \gamma_{ns}(\overline{G}) \leq (p - 2)^2. \quad \dots (23)$$

#### ACKNOWLEDGEMENT

Thanks are due to the referees for their valuable comments and suggestions.

#### REFERENCES

1. G. Chartrand and L. Lesniak, *Graphs and Digraphs*, Chapman and Hall, Madras. (1996).
2. E. J. Cockayne, Domination of undirected graphs-A survey. In *Theory and Applications of Graphs*. Lecture Notes in Mathematics 642, Springer - Verlag (1978) 141-47.
3. E. J. Cockayne and S. T. Hedetniemi, Towards a theory of domination in graphs. *Networks* 7 (1977) 247-61.
4. F. Harary, *Graph Theory*. Addison-Wesley, Reading Mass. (1969).
5. V. R. Kulli and B. Janakiram, The split domination number of a graph. *Graph Theory Notes of New York*, New York Academy of Sciences (1997) XXXII, 16-19.
6. E. Sampathkumar and H. B. Walikar, *J Math. Phys. Sci.* 13 (1979) 607-13.