

THE ORDER STRUCTURE OF WALLMAN NEARNESSSES

K. C. CHATTOPADHYAY AND SUSHAMA GUIN*

Department of Mathematics, University of Burdwan, Burdwan 713104
West Bengal

(Received 28 September 1992; after revision 26 May 1993;
accepted 30 August 1993)

Reed⁸ introduced Wallman nearnesses. In this article we investigate the order structure of the set $(\mathcal{M}_W(X), \subset)$ of all Wallman nearnesses compatible with a given T_1 -space X . It is well-known that the set $(\mathcal{M}(X), \subset)$ of all Lodato nearnesses compactible with the space X is a complete lattice. We describe explicitly the least and largest elements of $(\mathcal{M}_W(X), \subset)$ and prove that it is also a complete lattice. Though $\mathcal{M}_W(X) \subset \mathcal{M}(X)$, an example is given to show that $(\mathcal{M}_W(X), \subset)$ is not a sublattice of the lattice $(\mathcal{M}(X), \subset)$.

1. INTRODUCTION

In order to introduce Wallman type (W-T) extensions, Reed⁸ defined Wallman nearnesses. She proved that there is a 1-1 match between the set $\mathcal{M}_W(X)$ of all compatible Wallman nearnesses on a T_1 -space X and the set $\mathcal{E}_W(X)$ of all W-T extensions of X . Reed proved that all T_2 compactifications, in particular all Wallman-Frink compactifications, classical Wallman compactifications of T_1 spaces and one-point compactifications of T_1 spaces are W-T extensions. Recently present authors⁴ introduced a concept of generalised Wallman type compactifications and proved that they are also W-T extensions. Thus there is a large class of well-known compactifications belonging to the class of W-T extensions. Since for a T_1 -space X there is a 1-1 correspondence between $\mathcal{M}_W(X)$ and $\mathcal{E}_W(X)$, the order structure of $(\mathcal{M}_W(X), \subset)$ has some bearing on that of $(\mathcal{E}_W(X), \leq)$.

In this article, the investigation of the order structure of $(\mathcal{M}_W(X), \subset)$ for a T_1 -space X has just been begun which may be thought to be far from being completed. It is well-known that the set $(\mathcal{M}(X), \subset)$ of all Lodato nearnesses

*The research of the second author is supported by CSIR, New Delhi.

compatible with the space X is a complete lattice. We describe explicitly the least and the largest elements of $(\mathcal{M}_w(X), \subset)$ and prove that it is also a complete lattice. Though $\mathcal{M}_w(X) \subset \mathcal{M}(X)$, an example is given to show that $(\mathcal{M}_w(X), \subset)$ is not a sublattice of the lattice $(\mathcal{M}(X), \subset)$.

Preliminaries 1.1 — In what follows, in this section, we collect some definitions and results without proof from works of different authors which will be used in this article.

Theorem 1.2 — If in a partially ordered set every nonempty subset that has an upper bound has a least upper bound, then every nonempty subset which has a lower bound has a greatest lower bound. (For proof see Thron¹⁰).

Definition 1.3 — A ‘grill’ \mathcal{G} on X is a family of subsets of X satisfying : $A \in \mathcal{G}$ and $A \subset B \Rightarrow B \in \mathcal{G}$, $A \cup B \in \mathcal{G} \Rightarrow A \in \mathcal{G}$ or $B \in \mathcal{G}$ and $\emptyset \notin \mathcal{G}$.

Another way to look at grills is that they are the union of ultrafilters. The collection of all grills on X will be denoted by $\Gamma(X)$ (For more details on these concepts we refer to Thron⁹).

A grill \mathcal{G} on a topological space X is called a c -grill if

$$\forall A \subset X, \bar{A} \in \mathcal{G} \Rightarrow A \in \mathcal{G}.$$

For each point x of a topological space X , set,

$$\mathcal{G}_x = \{ A \subset X : x \in \bar{A} \}.$$

One can easily check that \mathcal{G}_x is a c -grill on X for each $x \in X$ and \mathcal{G}_x is called the ‘adherence grill’ of the point x .

Definition 1.4 — A ‘closed filter’ on a topological space X shall be a filter on X having a base consisting of closed sets only. An ‘ultraclosed filter’ on X shall be maximal closed filter on X . Using Zorn’s lemma it can be proved that every closed filter is contained in an ultraclosed filter.

Proposition 1.5 — If F is a closed set in X and F intersects each member of a base of an ultraclosed filter \mathcal{U} , consisting of only closed sets, then $F \in \mathcal{U}$.

For a topological space X we shall use the notation

$$\Psi(X) = \{ \mathcal{U} : \mathcal{U} \text{ is an ultraclosed filter on } X \}.$$

We note that if X is a T_1 -space then for every x in X , $\mathcal{U}_x = \{ A : x \in A \}$ is an ultraclosed filter on X and each such ultraclosed filter is called a principal ultraclosed filter. An ultraclosed filter is known as a nonprincipal (n.p.) ultraclosed filter if it is not a principal ultraclosed filter.

Results concerning closed filters may be found in Chattopadhyay¹ and Thorn¹⁰.

Definition 1.6 — A (basic) proximity π on X is a symmetric binary relation on the power set $\mathcal{P}(X)$ of X satisfying the following conditions :

$$P_1 : A \cap B \neq \phi \Rightarrow (A, B) \in \pi$$

$$P_2 : (A, B \cup C) \in \pi \Leftrightarrow (A, B) \in \pi \text{ or } (A, C) \in \pi$$

$$P_3 : (X, \phi) \notin \pi.$$

A (basic) proximity π on X induces a closure operator

$$c_\pi(A) = \{x \in X : (\{x\}, A) \in \pi\}.$$

A (basic) proximity π is called a LO-(Lodato) proximity if $(c_\pi(A), c_\pi(B)) \in \pi \Rightarrow (A, B) \in \pi$. A grill \mathcal{G} on X is called a π -clan if

$$(A, B) \in \pi \forall \{A, B\} \subset \mathcal{G}.$$

Definition 1.7 — A (basic) nearness ν on X is a collection of families $\mathcal{A} \subset \mathcal{P}(X)$ satisfying :

$$N_1 : \bigcap \mathcal{A} \neq \phi \Rightarrow \mathcal{A} \in \nu$$

$$N_2 : \mathcal{A} \in \nu \Rightarrow \phi \notin \nu$$

$$N_3 : \mathcal{A} \in \nu \Rightarrow \mathcal{B} \in \nu \text{ if } \mathcal{B} \text{ satisfies } B \in \mathcal{B} \Rightarrow \exists A \in \mathcal{A}, B \supset A$$

$$N_4 : \mathcal{A} \notin \nu, \mathcal{B} \notin \nu \Rightarrow \{A \cup B : A \in \mathcal{A}, B \in \mathcal{B}\} \notin \nu.$$

A (basic) nearness ν induces a (basic) proximity

$$\pi_\nu = \{(A, B) : \{A, B\} \in \nu\}.$$

A (basic) nearnesses induces a closure operator

$$c_\nu(A) = \{x : \{x\}, A \in \nu\}.$$

The (basic) nearness ν is called a LO-(Lodato) nearness (or simply a nearness due to Herrlich⁷ if

$$\{c_\nu(A) : A \in \mathcal{A}\} \in \nu \Rightarrow \mathcal{A} \in \nu.$$

A grill \mathcal{G} on X is a ν -clan if $\mathcal{G} \in \nu$. Thus $\nu \cap \Gamma(X)$ is the set of all ν -clans. A maximal family in ν is called a ν -cluster. Every ν -cluster is a grill and is maximal ν -clan. A nearness is called cluster generated (or concerte) if $\mathcal{A} \in \nu \Rightarrow \mathcal{A} \subset \mathcal{G}$ for some ν -cluster \mathcal{G} . The collection of all ν -cluster is denoted by X^ν .

Details on proximities and nearnesses may be found in Gagrat and Thron⁵, Herrlich⁶ and Thron⁹. Now we state Theorem 2.7 of Gagrat and Thron⁵ which will frequently be used in this article.

Theorem 1.8 — Let $\gamma = \{\mathcal{G}_i : i \in I\}$ be a family of grills on X with the property that for every $x \in X$ there exists an i such that $\{x\} \in \mathcal{G}_i$. Then ν_γ , defined by $\mathcal{A} \in \nu_\gamma$ if and only if $\mathcal{A} \subset \mathcal{G}_i$ for some $i \in I$, is a basic nearness on X . If in

addition no two members of γ are comparable then ν_γ is cluster generated and $X^{\nu_\gamma} = \gamma$.

Let X be a topological space. A grill \mathcal{G} on X is called a 'Wallman grill' if there exists an ultraclosed filter \mathcal{U} on X such that $\mathcal{U} \subset \mathcal{G}$. The set of all Wallman grills on X shall be denoted by $\Gamma^W(X)$. Thus

$$\Gamma^W(X) = \{\mathcal{G} \in \Gamma(X) : \exists \mathcal{U} \in \Psi(X), \mathcal{U} \subset \mathcal{G}\}.$$

Henceforth all spaces are assumed to be T_1 unless stated otherwise. Let π be a compatible LO -proximity on a space X ; that is, $c_\pi(A) = \bar{A}$ for every $A \subset X$ where \bar{A} denotes the closure of A in X . A π -clan \mathcal{G} is called a Wallman π -clan if $\mathcal{G} \in \Gamma^W(X)$. One can easily check that each adherence grill \mathcal{G}_x is a Wallman π -clan. Thus in view of Theorem 1.8, the set of all Wallman π -clans generates a nearness which is denoted by $\nu_W(\pi)$. Thus

$$\nu_W(\pi) = \{\mathcal{A} \subset \mathcal{P}(X) : \mathcal{A} \subset \mathcal{G} \text{ for some wallman } \pi\text{-clan } \mathcal{G}\}.$$

One can easily check that $\nu_W(\pi)$ is a compatible LO -nearness, that is, $c_{\nu_W(\pi)}(A) = \bar{A}$ for all $A \subset X$. A nearness of the form $\nu_W(\pi)$ is called a 'Wallman nearness' on X . Thus a Wallman nearness on X is a nearness generated by the set of all Wallman π -clans for some compatible LO -proximity π on X .

Proposition 1.9 — A compatible LO -nearness ν on a space X is a Wallman nearness if and only if $\nu = \nu_W(\pi_\nu)$. (For the proof see Reed⁸).

We conclude this section by collecting a few more known definitions and results with exact references for they will frequently be used in the rest of the paper.

Definition 1.10 (Chattopadhyay and Guin³, Definition 2.1) — A family of grills

$$\Gamma_1 = \{\mathcal{G}_i : i \in I\} \subset \Gamma(X)$$

is called a binary collection with respect to Γ_2 (a second family in $\Gamma(X)$) if the following holds :

If $\mathcal{G} \in \Gamma_2$ has the property that each two-element subfamily \mathcal{A} of \mathcal{G} implies the existence of $i(\mathcal{A}) \in I$ such that $\mathcal{A} \subset \mathcal{G}_{i(\mathcal{A})} \in \Gamma_1$ then there exists a $j \in I$ such that $\mathcal{G} \subset \mathcal{G}_j \in \Gamma_1$.

In what follows we give a simple but useful characterization of Wallman nearnesses.

Theorem 1.11³ — Let ν be a compatible cluster generated LO -nearness on a space X . Then ν is a Wallman nearness if and only if $X^\nu \subset \Gamma^W(X)$ and X^ν is a binary collection with respect to $\Gamma^W(X)$.

Definition 1.12³ — For a topological space X and $\mathcal{A} \subset \mathcal{P}(X)$

$$b(\mathcal{A}) = \{A \subset X : \bar{A} \in \mathcal{A}\}$$

then following properties of the operator 'b' can be easily verified.

Theorem 1.13³ — For a topological space X , the following results hold.

- (i) $\mathcal{F} \subset b(\mathcal{F})$ for every filter \mathcal{F} on X .
- (ii) If $\mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_n$ are filters on X then

$$b(\mathcal{F}_1 \cup \mathcal{F}_2 \cup \dots \cup \mathcal{F}_n) = b(\mathcal{F}_1) \cup b(\mathcal{F}_2) \cup \dots \cup b(\mathcal{F}_n).$$

- (iii) $b(\mathcal{U})$ is a c -grill whenever \mathcal{U} is an ultrafilter or ultraclosed filter on X .
- (iv) $b(\mathcal{U}_x) = \mathcal{G}_x$ for every principal ultrafilter \mathcal{U}_x on X .
- (v) If \mathcal{U} is an ultrafilter and \mathcal{V} is an ultraclosed filter on X such that $\mathcal{V} \subset \mathcal{U}$ then $\mathcal{U} \subset b(\mathcal{V})$.
- (vi) Let $\mathcal{V}_1, \mathcal{V}_2$ be two ultraclosed filters on X . Then

$$\mathcal{V}_1 \subset b(\mathcal{V}_2) \Leftrightarrow \mathcal{V}_1 = \mathcal{V}_2 \Leftrightarrow b(\mathcal{V}_1) \subset b(\mathcal{V}_2).$$

- (vii) Let $\mathcal{U}, \mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n$ be ultraclosed filters on X such that $\mathcal{U} \subset b(\mathcal{U}_1 \cup \mathcal{U}_2 \cup \dots \cup \mathcal{U}_n)$ then $\mathcal{U} = \mathcal{U}_i$ for some $i = 1, 2, \dots, n$.

Remark 1.14 : Though more general results can be proved than those stated in the above theorem, in view of our purpose we do not wish to do so.

Proposition 1.15³ — Let \mathcal{A} be a family of subsets of a topological space X such that $\overline{\mathcal{A}}$ has the finite intersection property (F.I.P.). Then there exists an ultraclosed filter \mathcal{U} on X such that $\mathcal{A} \subset b(\mathcal{U})$ where $\overline{\mathcal{A}} = \{\overline{A} : A \in \mathcal{A}\}$.

Proposition 1.16³ — Let ν be a compatible LO -nearness on a space X and \mathcal{U} be an ultraclosed filter on X . Then $b(\mathcal{U})$ is a Wallman π_ν -clan.

Above results immediately yield the following

Theorem 1.17³ — Let ν be a compatible Wallman nearness on a space X . Then

- (i) $b(\mathcal{U}) \in \nu$ for every $\mathcal{U} \in \Psi(X)$.
- (ii) $\mathcal{A} \in \nu$ if $\mathcal{A} \subset \mathcal{P}(X)$ and $\overline{\mathcal{A}}$ has the F.I.P.
- (iii) $b(\mathcal{F}) \in \nu$ for all filters \mathcal{F} on X .
- (iv) $\mathcal{F} \in \nu$ for all filters \mathcal{F} on X .

Theorem 1.18³ — Let X be a space and ν_0 be the elementary nearness on the space X , that is

$$\nu_0 = \{\mathcal{A} \subset \mathcal{P}(X) : \bigcap \overline{\mathcal{A}} \neq \phi\}.$$

Then following are equivalent :

- (i) X is compact.
- (ii) ν_0 is a compatible Wallman nearness on X .

2. THE ORDER STRUCTURE OF WALLMAN NEARNESSSES

This section contains our results concerning the order structure of the partially ordered set $(\mathcal{M}_w(X), \subset)$. Let $\mathcal{M}(X)$ be the set of all *LO*-nearnesses compatible with the space X . Note that $\mathcal{M}_w(X) \subset \mathcal{M}(X)$. Though it is well-known from Herrlich⁶ that $(\mathcal{M}(X), \subset)$ is a complete lattice, for the sake of completeness below we sketch a proof of it.

Note that $v_0 = \{\mathcal{A} \subset \mathcal{P}(X) : \bigcap \bar{\mathcal{A}} = \emptyset\}$ is the smallest element of $(\mathcal{M}(X), \subset)$.

Also if $\{v_i : i \in I\}$ is a nonempty subfamily of $\mathcal{M}(X)$ then it is immediate that

$$v = \bigcup \{v_i : i \in I\}$$

is an element of $\mathcal{M}(X)$ and hence v is the least upper bound of $\{v_i : i \in I\}$ in the partially ordered set $(\mathcal{M}(X), \subset)$ and hence in view of Theorem 1.2, we conclude that $(\mathcal{M}(X), \subset)$ is a complete lattice.

In order to prove that $(\mathcal{M}_w(X), \subset)$ is a complete lattice we need the following two lemmas.

Lemma 2.1 — A cluster generated (basic) nearness v on a set X is a *LO*-nearness if and only if each v -cluster is a *c*-grill on (X, c_v) .

PROOF : If v is a cluster generated *LO*-nearness and \mathcal{G} is a v -cluster then $c_v(A) \in \mathcal{G}$ implies that $\{c_v(A)\} \cup \{c_v(B) : B \in \mathcal{G}\} \in v$ and since v is a *LO*-nearness, $\{A\} \cup \mathcal{G} \in v$ and hence by the maximality of \mathcal{G} , $A \in \mathcal{G}$.

Conversely suppose that v is a cluster generated (basic) nearness such that each v -cluster is a *c*-grill on (X, c_v) . Let $\{c_v(A) : A \in \mathcal{A}\} \in v$ for some $\mathcal{A} \subset \mathcal{P}(X)$. Since v is cluster generated, there exists a v -cluster \mathcal{G} such that $\{c_v(A) : A \in \mathcal{A}\} \subset \mathcal{G}$. Since \mathcal{G} is a *c*-grill, $\mathcal{A} \subset \mathcal{G}$ and hence $\mathcal{A} \in v$. Thus v is a *LO*-nearness on X .

Lemma 2.2 — Let X be a topological space.

Set

$$\gamma_1 = \{b(\mathcal{U}) : \mathcal{U} \in \Psi(X)\}.$$

Then

$$v_{\gamma_1} = \{\mathcal{A} \subset \mathcal{P}(X) : \mathcal{A} \subset \mathcal{G} \text{ for some } \mathcal{G} \in \gamma_1\}$$

is a compatible Wallman nearness on X such that $X^{v_{\gamma_1}} = \gamma_1$.

PROOF : Obviously for $x \in X$, \mathcal{U}_x is an ultraclosed filter and $\{x\} \in b(\mathcal{U}_x)$. Also in view of Theorem 1.13 (iii) and (vi) it follows that if \mathcal{U}, \mathcal{V} are ultraclosed filters then $b(\mathcal{U})$ is a *c*-grill and $b(\mathcal{U}) \subset b(\mathcal{V})$ implies that $\mathcal{U} = \mathcal{V}$.

Taking into account these facts we conclude, by Theorem 1.8 and Lemma 2.1, that ν_{γ_1} is a cluster generated *LO*-nearness compatible with the space X and $X^{\nu_{\gamma_1}} = \gamma_1$.

To show that ν_{γ_1} is a Wallman nearness we need to check that $\gamma_1 \subset \Gamma^W(X)$ and γ_1 is binary with respect to $\Gamma^W(X)$ (Theorem 1.11). Since each member of γ_1 is of the form $b(\mathcal{U})$ for some ultraclosed filter \mathcal{U} on X it follows by Theorem 1.13 (i) that $\gamma_1 \subset \Gamma^W(X)$.

Let $\mathcal{G} \in \Gamma^W(X)$ such that each two-element subfamily of \mathcal{G} is contained in some element of γ_1 . Choose $\mathcal{U}_0 \in \Psi(X)$ such that $\mathcal{U}_0 \subset \mathcal{G}$. Let $A \in \mathcal{G}$ and F be any arbitrary closed set in \mathcal{U}_0 . Then since $\{A, F\}$ is contained in some member $b(\mathcal{U})$ of γ_1 it follows that $\bar{A} \cap F \neq \emptyset$. Since the closed set F has been chosen arbitrarily in \mathcal{U}_0 and \mathcal{U}_0 is ultraclosed, in view of Proposition 1.5, $\bar{A} \in \mathcal{U}_0$ and hence $A \in b(\mathcal{U}_0)$, consequently $\mathcal{G} \subset b(\mathcal{U}_0)$. Thus γ_1 is a binary collection with respect to $\Gamma^W(X)$.

Theorem 2.3 — Let X be a space. Then $(\mathcal{M}_W(X), \subset)$ is a complete lattice with

$$\{\mathcal{A} \subset \mathcal{P}(X) : \mathcal{A} \subset b(\mathcal{U}), \mathcal{U} \in \Psi(X)\}$$

as the least element of $(\mathcal{M}_W(X), \subset)$.

PROOF : Denote by

$$\gamma_1 = \{b(\mathcal{U}) : \mathcal{U} \in \Psi(X)\}.$$

Then by the above lemma ν_{γ_1} is an element of $\mathcal{M}_W(X)$. If $\nu \in \mathcal{M}_W(X)$ then $b(\mathcal{U}) \in \nu$ for all $\mathcal{U} \in \Psi(X)$ (see Theorem 1.17(i)). Hence ν_{γ_1} is the least element of $(\mathcal{M}_W(X), \subset)$.

Let $\{\nu_i : i \in I\}$ be a nonempty subfamily of $\mathcal{M}_W(X)$. Let $\nu = \bigcup \{\nu_i : i \in I\}$. Since $\{\nu_i : i \in I\} \subset \mathcal{M}(X)$ according to the discussion at the beginning of the section

$$\nu = \bigcup \{\nu_i : i \in I\} \in \mathcal{M}(X).$$

In general, ν need not be a Wallman nearness (see Example 2.6). Let us set

$$\nu_L = \nu_W(\pi_\nu).$$

In view of proposition 1.9, ν_L belongs to $\mathcal{M}_W(X)$. We shall show that ν_L is the l.u.b. of $\{\nu_i : i \in I\}$ in $(\mathcal{M}_W(X), \subset)$.

Let $i \in I$ and $\mathcal{A} \in \nu_i$. Then there exists a Wallman π_{ν_i} -clan \mathcal{G} such that $\mathcal{A} \subset \mathcal{G}$. Since $\nu_i \subset \nu$, $\pi_{\nu_i} \subset \pi_\nu$ and hence \mathcal{G} is a Wallman π_ν -clan, consequently $\mathcal{A} \in \nu_L$. Thus ν_L is an upper bound of $\{\nu_i : i \in I\}$ in $(\mathcal{M}_W(X), \subset)$.

Let v^* be an element of $\mathcal{M}_W(X)$ such that $v_i \subset v^*$ for all $i \in I$. Then $v \subset v^*$ and hence $\pi_v \subset \pi_{v^*}$. Thus any Wallman π_v -clan is a Wallman π_{v^*} -clan. Hence

$$v_L = v_W(\pi_v) \subset v_W(\pi_{v^*}) = v^*.$$

Thus in view of Theorem 1.2, we conclude that $(\mathcal{M}_W(X), \subset)$ is a complete lattice.

Remark 2.4 : We have seen that $(\mathcal{M}(X), \subset)$ and $(\mathcal{M}_W(X), \subset)$ are both complete lattices whenever X is a space. Also $\mathcal{M}_W(X) \subset \mathcal{M}(X)$. We shall give an example to show that $(\mathcal{M}_W(X), \subset)$ is not a sublattice of $(\mathcal{M}(X), \subset)$. To accomplish it we begin with the following example.

Example 2.5 — Existence of uncountably many n.p. ultraclosed filters on a noncompact space.

Let X be an uncountable set with the co-countable topology. From a well-known result of set theory we assert that there exists a partition $\{X_x : x \in X\}$ of X ,

i.e., $X = \bigcup \{X_x : x \in X\}$ and $X_x \cap X_y = \emptyset$ if $x \neq y$ satisfying the additional property

$$|X_x| = |X| \quad \forall x \in X$$

where $|X|$ denotes the cardinal number of the set X .

Now let us choose $F_x \subset X_x \quad \forall x \in X$ such that $|F_x| = \aleph_0$. Since each F_x is countable, it can be expressed as

$$F_x = \{Y_{x,1}, Y_{x,2}, Y_{x,3}, \dots\} \quad \forall x \in X.$$

Now set

$$B_{x,n} = \{Y_{x,n}, Y_{x,n+1}, Y_{x,n+2}, \dots\} \quad \forall n \in \mathbb{N}, x \in X.$$

and

$$\mathcal{B}_x = \{B_{x,1}, B_{x,2}, B_{x,3}, \dots\} \quad \forall x \in X.$$

Then for each $x \in X$, \mathcal{B}_x is a family of closed sets with the F.I.P. Hence there exists an ultraclosed filter $\mathcal{U}(\mathcal{B}_x)$ such that $\mathcal{B}_x \subset \mathcal{U}(\mathcal{B}_x)$. Since $\bigcap \mathcal{B}_x = \emptyset$, $\mathcal{U}(\mathcal{B}_x)$ is n.p.

Finally note that if $x, y \in X$ and $x \neq y$ then $X_x \cap X_y = \emptyset$. Then $\mathcal{U}(\mathcal{B}_x) \neq \mathcal{U}(\mathcal{B}_y)$ for $X_x \in \mathcal{U}(\mathcal{B}_x)$ and $X_y \in \mathcal{U}(\mathcal{B}_y)$. Thus $|\mathcal{U}(\mathcal{B}_x) : x \in X| = |X|$.

Example 2.6 — Union of Wallman nearnesses on a space X may not be a Wallman nearness.

Let X be a concompact space and Φ be a set of n.p. ultraclosed filters on X such that Φ contains at least three elements (such spaces exist), see the example given above. Let ξ be a collection of finite proper subsets of Φ such that ξ contains each two-element subset of Φ .

Now set

$$\gamma_{\mathcal{A}} = \{b(\mathcal{U}) : \mathcal{U} \in \Psi(X) - \mathcal{A}\} \cup \{b(\bigcup \mathcal{A})\} \quad \forall \mathcal{A} \in \xi.$$

In view of Theorems 1.8, 1.13 (iii) and Lemma 2.1, it immediately follows that $\nu_{\gamma_{\mathcal{A}}}$ is a cluster generated *LO*-nearness with $\gamma_{\mathcal{A}}$ as the set of $\nu_{\gamma_{\mathcal{A}}}$ -clusters for each $\mathcal{A} \in \xi$.

Let $\mathcal{A}_0 \in \xi$. Since \mathcal{A}_0 is a finite subset of Φ , \mathcal{A}_0 can be expressed as $\mathcal{A}_0 = \{\mathcal{U}_1, \mathcal{U}_2, \dots, \mathcal{U}_n\}$. We now show that $\nu_{\gamma_{\mathcal{A}_0}}$ is a Wallman nearness. Clearly $\gamma_{\mathcal{A}_0} \subset \Gamma^W(X)$. Thus to check that $\nu_{\gamma_{\mathcal{A}_0}}$ is a Wallman nearness it is sufficient to check that $\gamma_{\mathcal{A}_0}$ is a binary collection with respect to $\Gamma^W(X)$.

Let $\mathcal{G} \in \Gamma^W(X)$ such that each two-element subfamily of \mathcal{G} is contained in some element of $\gamma_{\mathcal{A}_0}$. Since $\mathcal{G} \in \Gamma^W(X)$, \mathcal{G} contains an ultraclosed filter \mathcal{U}_0 on X .

Case 1 — $\mathcal{U}_0 \in \Psi(X) - \mathcal{A}_0$. Then we claim that $\mathcal{G} \subset b(\mathcal{U}_0)$. If not, by Proposition 1.5, there is a closed set $F \in \mathcal{U}_0$ and $A \in \mathcal{G}$ such that $\bar{A} \cap F = \phi$. Also since $\mathcal{U}_0 \notin \mathcal{A}_0$, there are closed sets F_1, F_2, \dots, F_n in \mathcal{U}_0 such that $F_k \notin \mathcal{U}_k \quad \forall k = 1, 2, \dots, n$. Let $B = F \cap F_1 \cap F_2 \cap \dots \cap F_n$. Then $B \in \mathcal{U}_0 \subset \mathcal{G}$. Thus $\{A, B\} \subset \mathcal{G}$. Since $\bar{A} \cap B = \phi$, $\{A, B\}$ is not contained in any element of $\gamma_{\mathcal{A}_0}$ -a contradiction to the fact that each two-element subfamily of \mathcal{G} is contained in an element of $\gamma_{\mathcal{A}_0}$.

Case 2 — $\mathcal{U}_0 \in \mathcal{A}_0$. Then we claim that $\mathcal{G} \subset b(\bigcup \mathcal{A}_0)$. If not, there is $A \in \mathcal{G}$ such that $\bar{A} \notin \bigcup \mathcal{A}_0$ and hence in particular $\bar{A} \notin \mathcal{U}_0$. Then there is a closed set $B \in \mathcal{U}_0$ such that $\bar{A} \cap B = \phi$. Clearly $\{A, B\} \not\subset b(\bigcup \mathcal{A}_0)$ because $A \notin b(\bigcup \mathcal{A}_0)$. Also since $\bar{A} \cap B = \phi$, $\{A, B\} \not\subset b(\mathcal{U})$ for all $\mathcal{U} \in \Psi(X) - \mathcal{A}_0$. Note that $\{A, B\}$ is a two-element subfamily of \mathcal{G} which is not contained in any element of $\gamma_{\mathcal{A}_0}$ -a contradiction.

Since \mathcal{A}_0 has been chosen arbitrarily in ξ , $\nu_{\gamma_{\mathcal{A}}}$ is a Wallman nearness for each $\mathcal{A} \in \xi$.

Set

$$\gamma = \{b(\mathcal{U}) : \mathcal{U} \in \Psi(X) - \bigcup \xi\} \cup \{b(\bigcup \mathcal{A}) : \mathcal{A} \in \xi\}.$$

Note that $\gamma \subset \bigcup \{\gamma_{\mathcal{A}} : \mathcal{A} \in \xi\}$ and each element of $\bigcup \{\gamma_{\mathcal{A}} : \mathcal{A} \in \xi\}$ is contained in some element of γ . Thus

$$\nu_{\gamma} = \bigcup \{\nu_{\gamma_{\mathcal{A}}} : \mathcal{A} \in \xi\}$$

It can be easily proved that ν_{γ} is a cluster generated *LO*-nearness on X with γ as the set of ν_{γ} -clusters. In fact each $\nu_{\gamma_{\mathcal{A}}}$ and ν_{γ} are compatible with the space X .

Now we show that ν_γ fails to be a Wallman nearness. Since each n.p. ultraclosed filter is contained in a n.p. ultrafilter, we can choose a n.p. ultrafilter \mathcal{U}^* for each $\mathcal{U} \in \Phi$ such that $\mathcal{U} \subset \mathcal{U}^*$. Set

$$\mathcal{G}^* = \bigcup \{ \mathcal{U} : \mathcal{U} \in \Phi \}.$$

By Theorem 1.13 (v), $\mathcal{U}^* \subset b(\mathcal{U})$ for all $\mathcal{U} \in \Phi$. Note that $\mathcal{G}^* \subset \Gamma^w(X)$ and each two-element subfamily of \mathcal{G}^* is contained in some element of $\{b(\bigcup \mathcal{A}) : \mathcal{A} \in \xi\}$ but in view of 1.13(vi), (vii), \mathcal{G}^* is not contained in any element of γ because $\mathcal{A} \neq \Phi$ for each $\mathcal{A} \in \xi$. Thus γ fails to be a binary collection with respect to $\Gamma^w(X)$. Since $\gamma = X^\gamma$, by Theorem 1.11, ν_γ is not a Wallman nearness.

Now as a corollary of the above example it follows that the l.u.b. of $\{\nu_{\gamma_{\mathcal{A}}} : \mathcal{A} \in \xi\}$ in $(\mathcal{M}_w(X), \subset)$ is different from $\nu_\gamma = \bigcup \{\nu_{\gamma_{\mathcal{A}}} : \mathcal{A} \in \xi\}$ which is the l.u.b. of $\{\nu_{\gamma_{\mathcal{A}}} : \mathcal{A} \in \xi\}$ in $(\mathcal{M}(X), \subset)$ (see the discussion at the beginning of this section). Thus $(\mathcal{M}_w(X), \subset)$ is not a sublattice of the lattice $(\mathcal{M}(X), \subset)$.

Following two theorems give an explicit description of the largest element in $(\mathcal{M}_w(X), \subset)$. In order to prove the theorems we need the following lemma.

Lemma 2.7 — A space X is compact if and only if each ultraclosed filter is principal.

PROOF : Suppose that X is compact. Let \mathcal{U} be an ultraclosed filter on X . Since X is compact, there exists an $x \in X$ such that $x \in \bar{A}$ for all $A \in \mathcal{U}$. Since \mathcal{U} is ultraclosed and X is T_1 , by Proposition 1.5 $\{x\} \in \mathcal{U}$ and hence $\mathcal{U} \subset \mathcal{U}_x$. Since \mathcal{U}_x is closed filter, by the maximality $\mathcal{U} = \mathcal{U}_x$.

Conversely suppose that X is not compact. Then there exists a nonempty family \mathcal{F} of closed sets in X with the finite intersection property such that $\bigcap \mathcal{F} = \phi$. Clearly \mathcal{F} is a subbase of a closed filter on X . Using Zorn's lemma one can prove that there exists an ultraclosed filter \mathcal{U} such that $\mathcal{F} \subset \mathcal{U}$. Since $\bigcap \mathcal{F} \supset \mu$ and $\bigcap \mathcal{F} = \phi$ it follows that $\bigcap \mathcal{U} = \phi$ and hence \mathcal{U} is nonprincipal.

Theorem 2.8 — If X is compact then the elementary nearness ν_0 is the only compatible Wallman nearness on X . In particular ν_0 is the largest element in $(\mathcal{M}_w(X), \subset)$.

PROOF : Let X be compact and ν be a compatible Wallman nearness on X . Since ν is a LO-nearness, $\mathcal{G}_x \in \nu$ for each $x \in X$. If $A \subset X$ and $\mathcal{G}_x \cup \{A\} \in \nu$ then $\{\{x\}, A\} \in \nu$ which implies that $x \in \bar{A}$, consequently $A \in \mathcal{G}_x$ and hence \mathcal{G}_x is a ν -cluster for each $x \in X$. Let \mathcal{G} be a ν -cluster. Then \mathcal{G} contains an ultraclosed filter on X . Since X is compact, in view of Lemma 2.7, there exists $x \in X$ such that

$\mathcal{U}_x \subset \mathcal{G}$ which implies, as above, $x \in \bar{A}$ for all $A \in \mathcal{G}$, consequently $\mathcal{G} \subset \mathcal{G}_x$ and hence $\mathcal{G} = \mathcal{G}_x$. Thus $\{\mathcal{G}_x : x \in X\}$ is the set of all ν -clusters, $\nu \subset \nu_0$. Since ν_0 is the smallest element in $(\mathcal{M}(X, \mathcal{C}), \nu = \nu_0)$. It is also known that ν_0 is a compatible Wallman nearness on X (Theorem 1.18). Thus ν_0 is the only element of $(\mathcal{M}_w(X, \mathcal{C}))$.

Theorem 2.9 — If X is a noncompact T_1 -space then

$$\nu_G = \{ \mathcal{A} \subset P(X) : \bigcap \bar{\mathcal{A}} \neq \emptyset \} \cup \{ \mathcal{A} \subset P(X) : A \in \mathcal{A} \Rightarrow |A| \geq \mathcal{H}_0 \}$$

is the largest element of $(\mathcal{M}_w(X, \mathcal{C}))$.

PROOF : Set

$$\gamma = \{ \mathcal{G}_x : x \in X \} \cup \{ \mathcal{G}^* \}$$

where $\mathcal{G}^* = \{ A \subset X : |A| \geq \mathcal{H}_0 \}$.

Then it is easily verified that $\nu_G = \nu_\gamma$. Clearly ν_G is a cluster generated LO-nearness compatible with the space X and $X^{\nu_G} = \gamma$.

Since X is T_1 , \mathcal{U}_x is ultraclosed filter for each $x \in X$. Also $\mathcal{U}_x \subset \mathcal{G}_x$ for each $x \in X$. Also since X is noncompact, in view of lemma 2.7, there is a non-principal ultraclosed filter \mathcal{U} on X . Clearly $\mathcal{U} \subset \mathcal{G}^*$. Thus $x^{\nu_\gamma} (= \gamma) \subset \Gamma^W(X)$.

We now prove that X^{ν_G} is binary with respect to $\Gamma^W(X)$. Let $\mathcal{G} \in \Gamma^W(X)$ such that each two-element subfamily of \mathcal{G} is contained in some element of X^{ν_G} . Choose $\mathcal{U} \in \Psi(X)$ such that $\mathcal{U} \subset \mathcal{G}$.

Case 1 — If $\mathcal{U} = \mathcal{U}_x$ for some $x \in X$ then $\mathcal{G} \subset \mathcal{G}_x$.

Case 2 — If \mathcal{U} is n.p. then we claim that $\mathcal{G} \subset \mathcal{G}^*$. If not, then there is $A \in \mathcal{G}$ such that $|A| < \mathcal{H}_0$. Since \mathcal{G} is a grill there is $x \in X$ such that $\{x\} \in \mathcal{G}$ nu $\{x\} \notin \mathcal{U}$ and hence there is a closed set $F \in \mathcal{U}$ such that $\{x\} \cap F = \emptyset$ (Proposition 1.5). Note that $\{\{x\}, F\} \subset \mathcal{G}$ but $\{\{x\}, F\}$ is not contained in any element of $X^{\nu_G} (= \gamma)$ — a contradiction to the fact that each two-element subfamily of \mathcal{G} is contained in some element of X^{ν_G} .

Thus ν_G is Wallman.

Next let ν be any element of $\mathcal{M}_w(X)$.

Let $\mathcal{G} \in X^\nu$. If $|A| \geq \mathcal{H}_0$ for each $A \in \mathcal{G}$

then $\mathcal{G} \subset \mathcal{G}^*$.

If \mathcal{G} contains a finite set then there exists $x \in X$ such that $\{x\} \in \mathcal{G}$. Then for each $A \in \mathcal{G}$, $\{\{x\}, A\} \in \nu$. Since ν is compatible with the space X , $x \in \bar{A}$ for each $A \in \mathcal{G}$ and hence $\mathcal{G} \subset \mathcal{G}_x$. Since ν is cluster generated, $\nu \subset \nu_G$.

Thus ν_G is the largest element of $(\mathcal{M}_W(X), \subset)$.

ACKNOWLEDGEMENT

Authors would like to thank the referee for useful suggestions.

REFERENCES

1. K. C. Chattopadhyay, *Pacific J. Math.* **118** (1985), 59-62.
2. K. C. Chattopadhyay, Sushma Guin and W. J. Thron, *Glasnik Matematički* **24** (1989), 115-23.
3. K. C. Chattopadhyay and Sushma Guin, *Bull. Alld. Math. Soc.* **2** (1987), 63-82.
4. K. C. Chattopadhyay and Sushma Guin, Compact Wallman type extensions through closed bases, (preprint).
5. M. S. Gagrat and W. J. Thron, *Trans. Amer. Math. Soc.* **208** (1975), 103-25.
6. H. Herrlich, *General Topology and Appl.* **4** (1974), 191-212.
7. H. Herrlich, Topological structures, *Mathematical Centre Tracts*, **52** (1974), 59-122.
8. Ellen E. Reed, *Pacific J. Math.* **95** (1981), 443-59.
9. W. J. Thron, *Math. Ann.* **206** (1973), 35-62.
10. W. J. Thron, *Topological Structures*, Holt, Rinehart and Winston, New York, 1966.
11. H. Wallman, *Ann. Math.* **39** (1938), 112-26.