

BILATERAL GENERATING RELATIONS FOR A FUNCTION DEFINED BY GENERALIZED RODRIGUES FORMULA

B. D. AGRAWAL AND J. P. CHAUBEY

Department of Mathematics, Banaras Hindu University, Varanasi 221005

(Received 21 May 1979; after revision 23 June 1980)

Several authors, including Srivastava and Panda (1975) have defined polynomial sets by means of Rodrigues formulae in different forms. In this paper our attempt is to unify all such polynomials and to derive bilateral generating relations for the polynomial set  $R_n^{(\alpha, \beta)}(x)$  defined by (1.1) below.

1. We define the polynomial set  $R_n(x)$  by means of the Rodrigues formula

$$\begin{aligned}
 R_n^{(\alpha, \beta)}[x; a, b, c, d; p, q, \gamma, \xi; w(x)] &\equiv R_n(x) \\
 &= \frac{(ax^p + b)^{-\alpha} (cx^q + d)^{-\beta}}{K_n w(x)} T_{k; \lambda}^n \left[ (ax^p + b)^{\alpha + \gamma n} (cx^q + d)^{\beta + \xi n} W(x) \right], \\
 & \qquad \qquad \qquad n = 0, 1, 2, \dots \dots (1.1)
 \end{aligned}$$

where

$$T_{k; \lambda} = x^k(\lambda + xD), \quad D \equiv \frac{d}{dx} \qquad \dots (1.2)$$

$K_n$  is some constant and  $w(x)$  is any general function of  $x$ , differentiable any number of times.

Some important and obvious connections of our polynomial set with those of the earlier workers are:

(1) with  $p = q = 1, K_n = n!, \lambda = 0, w(x) = \exp(-px^r), a = d = 1, b = c = 0, \gamma = -k$  and  $\alpha = a + kn$  (1.1) reduces to the polynomial set  $G_n^{(a)}(x, r, p, k)$ , defined by Srivastava and Singhal (1971).

(2)  $p = q = 1, K_n = n!, \lambda = 0, k = -1$ , reduces the polynomial set  $R_n(x)$  to  $S_n^{(\alpha, \beta)}[x; a, b, c, d; \gamma, \xi; w(x)]$ , defined by Srivastava and Panda (1975). Obviously, the polynomial set defined by Patil and Thakare (1977), too becomes a special case of (1.1).

Main results of the present paper consist of the generating relations:

$$\begin{aligned} & \sum_{n=0}^{\infty} R_{n+m}^{(\alpha-\gamma n, \beta-\xi n)} [x; a, b, c, d; p, q, \gamma, \xi; w(x)] \frac{t^n}{n!} \\ &= (1 - ktX')^{-\lambda/k} \frac{w\{x(1 - ktX')^{-1/k}\}}{w(x)} \left[ \frac{ax^p(1 - ktX')^{-p/k} + b}{ax^p + b} \right]^\alpha \\ & \quad \times \left[ \frac{cx^q(1 - ktX')^{-q/k} + d}{cx^q + d} \right]^\beta R_m^{(\alpha, \beta)} [x(1 - ktX')^{-1/k}; \\ & \quad a, b, c, d; p, q, \gamma, \xi; w(x(1 - ktX')^{-1/k})] \dots(1.3) \end{aligned}$$

where

$$X' = x^k(ax^p + b)^\gamma (cx^q + d)^\xi \dots(1.4)$$

and

$$\begin{aligned} & \sum_{n=0}^{\infty} R_n^{(\alpha-\gamma n, \beta-\xi n)} [x; a, b, c, d; p, q, \gamma, \xi; w(x)] \Phi_n(y)t^n \\ &= (1 - ktX')^{-\lambda/k} \frac{w\{x(1 - ktX')^{-1/k}\}}{w(x)} \left[ \frac{ax^p(1 - ktX')^{-p/k} + b}{ax^p + b} \right]^\alpha \\ & \quad \times \left[ \frac{cx^q(1 - ktX')^{-q/k} + d}{cx^q + d} \right]^\beta F[x(1 - ktX')^{-1/k}; X^n] \dots(1.5) \end{aligned}$$

where  $X'$  is given by (1.4),

$$\begin{aligned} X^n &= yt \left[ \frac{ax^p(1 - ktX')^{-p/k} + b}{ax^p + b} \right]^\gamma \left[ \frac{cx^q(1 - ktX')^{-q/k} + d}{cx^q + d} \right]^\xi \\ F[x; t] &= \sum_{n=0}^{\infty} \mu_n R_n^{(\alpha-\gamma n, \beta-\xi n)} [x; a, b, c, d; p, q, \gamma, \xi; w(x)] \frac{t^n}{n!} \dots(1.6) \end{aligned}$$

$\Phi_n(y)$  is a polynomial of degree  $n$  in  $y$  given by

$$\Phi_n(y) = \frac{1}{n!} \sum_{j=0}^n \binom{n}{j} \mu_j y^j$$

and  $\mu_j \neq 0$  are arbitrary constants.

§2. We now establish (1.3) from (1.1). We have

$$\sum_{n=0}^{\infty} R_{n+m}^{(\alpha-\gamma n+m, \beta-\xi n+m)} [x] \frac{t^n}{n!} =$$

$$= \frac{(ax^p + b)^{-\alpha + \gamma m} (cx^q + d)^{-\beta + \xi n}}{w(x)} \sum_{n=0}^{\infty} \frac{t^n}{n!} (ax^p + b)^{\gamma n} (cx^q + d)^{\xi n} \\ \times T_{k;\lambda}^m \{ (ax^p + b)^\alpha (cx^q + d)^\beta w(x) \}.$$

Using (1.1) and a property of  $T_{k;\lambda}$ , viz.

$$\exp(tT_{k;\lambda}) f(x) = (1 - kt x^k)^{-\lambda/k} f(x(1 - kt x^k)^{-1/k})$$

and after a little calculation we get the required result (1.3).

§3. For the proof of bilateral generating relation (1.5), let us start with (cf. Singhal and Srivastava 1972, p. 756, section 2)

$$\sum_{n=0}^{\infty} R_n^{(\alpha - \gamma n, \beta - \xi n)} [x] \Phi_n(y) t^n \\ = \sum_{j=0}^{\infty} \mu_j \frac{(yt)^j}{j!} \sum_{n=0}^{\infty} \frac{t^n}{n!} R_{n+j}^{(\alpha - \gamma n - \gamma j, \beta - \xi n - \xi j)} [x].$$

Now the result (1.5) would follow, if we interpret the above expression by (1.3) and (1.6) (see also Srivastva and Panda 1975, p. 312, Theorem 1).

§4. *Applications* — Since the polynomial set  $R_n(x)$  incorporates in itself several classical as well as other polynomials, a large variety of generating relations for the abovementioned polynomials may be obtained by assigning different values to the parameters in  $R_n^{(\alpha, \beta)}(x)$ .

REFERENCES

Patil, K. R., and Thakare, N. K. (1977). Bilateral generating function for a function defined by generalized Rodrigues formula. *Indian J. pure appl. Math.*, **8**, 425-29.  
 Singhal, J. P., and Srivastava, H. M. (1972). A class of bilateral generating functions for certain classical polynomials. *Pacific J. Math.*, **42**, 755-62.  
 Srivastava, H. M., and Panda, R. (1975). On the unified presentation of certain classical polynomials. *Boll. Un. Mat. Ital.* (4), **12**, 306-14.  
 Srivastava, H. M., and Singhal, J. P. (1971). A class of polynomials defined by generalized Rodrigues formula. *Ann. Mat. Pura Appl.* (4), **90**, 75-85.