

EXISTENCE OF POSITIVE SOLUTIONS FOR A CLASS OF QUASILINEAR SINGULAR ELLIPTIC SYSTEMS INVOLVING CAFFARELLI-KOHN-NIRENBERG EXPONENT WITH SIGN-CHANGING WEIGHT FUNCTIONS

Salah Boulaaras^{*,**}, Rafik Guefaïa^{***} and Tahar Bouali^{***}

**Department of Mathematics, College of Sciences and Arts, Al-Rass,
Qassim University, Kingdom of Saudi Arabia*

***Laboratory of Fundamental and Applied Mathematics of Oran (LMFAO),
University of Oran 1, Ahmed Benbella, Algeria*

****Department of Mathematics and Computer Science, Larbi Tebessi University 12002
Tebessa, Algeria*

e-mails: saleh_boulaares@yahoo.fr; salahboulaaras@yahoo.com; boualitahar@yahoo.fr

(Received 26 September 2017; accepted 11 January 2018)

The paper deals with the existence of weak positive solutions for a new class of quasilinear singular elliptic systems involving critical Caffarelli–Kohn–Nirenberg exponent with sign-changing weight functions using the method of sub-super solutions. Our results are natural extensions from the previous ones in [3].

Key words : Caffarelli–Kohn–Nirenberg exponents; elliptic system, sub-supersolution method, sign-changing.

1. INTRODUCTION

During the past few years, the treatise of positive solutions of singular partial differential equations or systems has been an extremely active research area. The singular nonlinear problems emerge naturally and they take a main role in the interdisciplinary field between analysis, biology, geometry, mathematical physics, elasticity, etc.

This article deals with the existence of positive solutions of the following boundary value problem

$$\begin{cases} -\operatorname{div} \left(|x|^{-ap} |\nabla u|^{p-2} \nabla u \right) = \lambda |x|^{-(a+1)p+c_1} [g(x) A(u) + f(v)], & \text{in } \Omega, \\ -\operatorname{div} \left(|x|^{-bq} |\nabla v|^{q-2} \nabla v \right) = \lambda |x|^{-(b+1)q+c_2} [g(x) B(v) + h(u)], & \text{in } \Omega, \\ u = v = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where Ω is a bounded domain in $\mathbb{R}^N$ and

$$1 < p, q < N, 0 \leq a \leq \frac{N-p}{N},$$

with

$$0 \leq b \leq \frac{N-q}{N}, c_1, c_2 \text{ are positive parameters.}$$

$g(x)$ is a C^1 sign-changing the weight function, that possibly negative nearby the boundary and f, h, A, B are C^1 nondecreasing functions satisfy

$$A(0) \geq 0, B(0) \geq 0.$$

The study of this kind of problems is motivated by its different applications, for example, population genetics, in fluid mechanics, Newtonian fluids, glaciology and flow through porous media (see for more detail [4, 9, 16, 21]).

On the other hand, there is an extensive practical background for quasilinear elliptic systems have. They are described in the multiplicative chemical reaction stimulated by the catalyst grains under variant temperature or constant, in the quasi-regular and quasi-conformal theory mappings in Riemannian manifolds with boundary, or in the description of many physical phenomena such as the pulses propagation in Kerr-like photorefractive media and birefringent optical fibers (see [19, 30]). Moreover, for additional results on elliptic problems, see ([1, 3, 6, 20, 24]). For the regular case, the quasilinear elliptic equation has been intensively studied by many authors where $c_1 = p, c_2 = q$ and $a = b = 0$, (see for example [2]). In the current work we concentrate on further extending the study in [5] for the quasilinear elliptic systems involving singularity. The extensions are nontrivial and challenging due to the singularity in the weights. Our approach is based on the method of sub and super solutions.

2. TECHNICAL ASSUMPTIONS AND AUXILIARY RESULTS

Let Ω be a bounded domain in $\mathbb{R}^N$ with its smooth boundary, $0 \in \Omega$ and $W_0^{1,p}(\Omega, |x|^{-ap})$ denote the completion of $C_0^\infty(\Omega)$ with the norm

$$\|u\| = \left(\int_{\Omega} |x|^{-ap} |\nabla u|^p dx \right)^{\frac{1}{p}}.$$

We consider the following nonlinear eigenvalue problem

$$\begin{cases} -\operatorname{div} \left(|x|^{-sr} |\nabla \phi|^{r-2} \nabla \phi \right) = \lambda |x|^{-(s+1)r+t} |\phi|^{r-2} \phi, x \in \Omega, \\ \phi = 0, x \in \partial\Omega. \end{cases} \quad (2.1)$$

For $t = c_1$ and $r = p, s = a$, we assume $\phi_{1,p}$ the eigenfunction corresponding to the first eigenvalue $\lambda_{1,p}$ of problem (2.1) where

$$\phi_{1,p} > 0 \text{ in } \Omega \text{ and } \|\phi_{1,p}\|_{\infty} = 1.$$

We consider the following assumptions similar to that in [26]:

For $s = b, r = q$, and $t = c_2$, we assume $\phi_{1,q}$ the eigenfunction corresponding to the first eigenvalue $\lambda_{1,q}$ of problem (2.1) where

$$\phi_{1,q} > 0 \text{ in } \Omega \text{ and } \|\phi_{1,q}\|_{\infty} = 1.$$

The maximum principle gives that

$$\frac{\partial \phi_r}{\partial n} < 0 \text{ on } \partial\Omega \text{ for } r \in \{p, q\},$$

where n is the outward normal. Then, there are positive constants m_0, δ and $\sigma_p, \sigma_q \in (0, 1)$ such that

$$\begin{cases} \lambda_{1,r} |x|^{-(s+1)r+t} \phi_{1,r}^r - |x|^{-sr} |\nabla \phi_{1,r}|^r \leq -m_0, x \in \overline{\Omega}_{\delta}. \\ \phi_{1,r} \geq \sigma_r, x \in \Omega \setminus \overline{\Omega}_{\delta}, \end{cases} \quad (2.2)$$

with

$$r \in \{p, q\}, s \in \{a, b\}, t \in \{c_1, c_2\} \text{ and } \overline{\Omega}_{\delta} = \{x \in \Omega : d(x, \partial\Omega) \leq \delta\}.$$

We also assume the unique solution $(\zeta_p(x), \zeta_q(x)) \in W_0^{1,p}(\Omega, |x|^{-ap}) \times W_0^{1,q}(\Omega, |x|^{-bq})$ of the following quasilinear singular system

$$\begin{cases} -\operatorname{div}(|x|^{-ap} |\zeta_p|^{p-2} \nabla \zeta_p) = |x|^{-(a+1)p+c_1}, x \in \Omega, \\ -\operatorname{div}(|x|^{-bq} |\zeta_q|^{q-2} \nabla \zeta_q) = |x|^{-(b+1)q+c_2}, x \in \Omega, \\ \zeta_p = \zeta_q = 0, x \in \partial\Omega, \end{cases} \quad (2.3)$$

where

$$\|\zeta_p\|_{\infty} = \mu_p \text{ and } \|\zeta_q\|_{\infty} = \mu_q.$$

Then, according [26], we have $\zeta_r > 0$ in and $\frac{\partial \phi_r}{\partial n} < 0$ on $\partial\Omega$ for $r \in \{p, q\}$.

Throughout this work, we consider that the weight function $g(x)$ hold negative values in $\overline{\Omega}_\delta$. However, it requires to be strictly positive in $\Omega \setminus \overline{\Omega}_\delta$. Precisely, we assume that there exist a positive constants β and η satisfy

$$g(x) \geq -\beta \text{ on } \overline{\Omega}_\delta \text{ and } g(x) \geq \eta \text{ on } \Omega \setminus \overline{\Omega}_\delta. \quad (2.4)$$

Let $s_0 \geq 0$ such that

$$\eta A(s) + f(s) > 0, \eta B(s) + h(s) > 0 \text{ for } s > s_0 \quad (2.5)$$

and

$$f_0 = \max\{0, -f(0)\}, h_0 = \max\{0, -h(0)\}. \quad (2.6)$$

3. MAIN RESULT

Putting

$$X = W_0^{1,p}(\Omega, |x|^{-ap}) \times W_0^{1,q}(\Omega, |x|^{-bq}).$$

We give the following definition of weak solution and sub-super solution of the problem (1.1):

Definition 1 — A pair of nonnegative functions $(\psi_1, \psi_2), (z_1, z_2)$ in X are called a weak sub-solution and supersolution of (1.1) if they satisfy: for $(\psi_1, \psi_2) = (z_1, z_2) = (0, 0)$ on $\partial\Omega$

$$\int_{\Omega} |x|^{-ap} |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \omega_1 dx \leq \lambda \int_{\Omega} \lambda |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)] \omega_1 dx,$$

$$\int_{\Omega} |x|^{-bq} |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \omega_2 dx \leq \lambda \int_{\Omega} |x|^{-(b+1)q+c_2} [g(x) B(\psi_2) + h(\psi_1)] \omega_2 dx$$

and

$$\int_{\Omega} |x|^{-ap} |\nabla z_1|^{p-2} \nabla z_1 \nabla \omega_1 dx \geq \lambda \int_{\Omega} |x|^{-(a+1)p+c_1} [g(x) A(z_1) + f(z_2)] \omega_1 dx,$$

$$\int_{\Omega} |x|^{-bq} |\nabla z_2|^{q-2} \nabla z_2 \nabla \omega_2 dx \geq \lambda \int_{\Omega} |x|^{-(b+1)q+c_2} [g(x) B(z_2) + h(z_1)] \omega_2 dx,$$

for all test functions

$$\omega_1(x) \in W_0^{1,p}(\Omega, |x|^{-ap})$$

and

$$\omega_2(x) \in W_0^{1,p}(\Omega, |x|^{-bq}),$$

with $\omega_1, \omega_2 \geq 0$. Then the following result holds:

Lemma 1 — (ref. [20]). Suppose there exist sub and super-solutions (ψ_1, ψ_2) and (z_1, z_2) respectively of (1.1) such that $(\psi_1, \psi_2) \leq (z_1, z_2)$. Then (1.1) has a weak solution (u, v) such that

$$(u, v) \in [(\psi_1, \psi_2), (z_1, z_2)].$$

In order to give the main result of this paper, we consider the following assumptions:

(A1) we have for every constant $K > 0$:

$$\lim_{s \rightarrow +\infty} \frac{f\left(K \left(h(s)^{\frac{1}{q-1}}\right)\right)}{s^{p-1}} = 0.$$

(A2)

$$\lim_{s \rightarrow +\infty} f(s) = \lim_{s \rightarrow +\infty} h(s) = +\infty.$$

(A3)

$$\lim_{s \rightarrow +\infty} \frac{A(s)}{s^{p-1}} = \lim_{s \rightarrow +\infty} \frac{B(s)}{s^{p-1}} = 0.$$

(A4) If $\alpha_p = \frac{p-1}{p} \sigma^{\frac{p}{p-1}}$, $\alpha_q = \frac{q-1}{q} \sigma^{\frac{q}{q-1}}$, and $\bar{\alpha} = \min\{\alpha_p, \alpha_q\}$ then there exists $\gamma > \frac{s_0}{\bar{\alpha}}$ such that

$$\begin{aligned} & \max \left\{ \frac{\gamma \lambda_{1,p}}{\eta A \left(\gamma^{\frac{1}{p-1}} \alpha_p \right) + f \left(\gamma^{\frac{1}{q-1}} \alpha_q \right)}, \frac{\gamma \lambda_{1,q}}{\eta B \left(\gamma^{\frac{1}{q-1}} \alpha_q \right) + h \left(\gamma^{\frac{1}{p-1}} \alpha_p \right)} \right\} \\ & < \min \left\{ \frac{m_0 \gamma}{\beta A \left(\gamma^{\frac{1}{p-1}} \right) + f_0}, \frac{m_0 \gamma}{\beta B \left(\gamma^{\frac{1}{q-1}} \right) + h_0} \right\}. \end{aligned}$$

We recall that m_0, σ_p and σ_q are introduced in relation (2.2) while s_0 is defined in (2.5). We now state our main result for the problem (1.1).

Theorem 1 — Suppose that (A1) – (A4) hold, then for every $\lambda \in [A, B]$, system (1.1) has at least one positive weak solution.

PROOF : Choose $r > 0$ such that $r \leq \min \left\{ |x|^{-(a+1)p+c_1}, |x|^{-(b+1)q+c_2} \right\}$ in $\overline{\Omega}_\delta$.

We take $\gamma > \frac{s_0}{\bar{\alpha}}$ as in hypothesis (A4). Define

$$A = \max \left\{ \frac{\gamma \lambda_{1,p}}{\eta A \left(\gamma^{\frac{1}{p-1}} \alpha_p \right) + f \left(\gamma^{\frac{1}{q-1}} \alpha_q \right)}, \frac{\gamma \lambda_{1,q}}{\eta B \left(\gamma^{\frac{1}{q-1}} \alpha_q \right) + h \left(\gamma^{\frac{1}{p-1}} \alpha_p \right)} \right\}$$

and

$$B = \min \left\{ \frac{m_0 \gamma}{\beta A \left(\gamma^{\frac{1}{p-1}} \right) + f_0}, \frac{m_0 \gamma}{\beta B \left(\gamma^{\frac{1}{q-1}} \right) + h_0} \right\}.$$

Setting

$$\psi_1 = (\gamma r)^{\frac{1}{p-1}} \frac{p-1}{p} \phi_{1,p}^{\frac{p}{p-1}} \text{ and } \psi_2 = (\gamma r)^{\frac{1}{q-1}} \frac{q-1}{q} \phi_{1,q}^{\frac{q}{q-1}}.$$

We will check that (ψ_1, ψ_2) is a sub-solution of (1.1) for $\lambda \in [A, B]$.

Indeed, let ω_1 with $\omega_1 \geq 0$ in. Then, it can be shown that

$$\left\{ \begin{aligned} \int_{\Omega} |x|^{-ap} |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \omega_1 dx &= \gamma r \int_{\Omega} |x|^{-ap} \phi_{1,p} |\nabla \phi_{1,p}|^{p-2} \nabla \phi_{1,p} \cdot \nabla \omega_1 dx \\ &= \gamma r \left\{ \int_{\Omega} |x|^{-ap} |\nabla \phi_{1,p}|^{p-2} \nabla \phi_{1,p} \cdot [\nabla (\phi_{1,p} \omega_1) - \nabla \phi_{1,p} \omega_1] dx \right\} \\ &= \gamma r \int_{\Omega} \left[\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right] \omega_1 dx. \end{aligned} \right. \quad (3.1)$$

Similarly we get □

$$\int_{\Omega} |x|^{-bq} |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \omega_2 dx = \gamma r \int_{\Omega} \left[\lambda_{1,q} |x|^{-(b+1)q+c_2} \phi_{1,q}^q - |x|^{-bq} |\nabla \phi_{1,q}|^q \right] \omega_2 dx. \quad (3.2)$$

Now, on $\overline{\Omega}_{\delta}$ by relation (2.2) we have

$$\gamma \left(\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right) \leq -m_0 \gamma.$$

Since $\lambda \leq B$ then we have

$$\lambda \leq \frac{m_0 \gamma}{\beta A \left(\gamma^{\frac{1}{p-1}} \right) + f_0},$$

thus

$$\left\{ \begin{aligned} \gamma \left(\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right) &\leq -m_0 \gamma \\ &\leq \lambda \left(-\beta A \left(\gamma^{\frac{1}{p-1}} \right) - f_0 \right) \\ &\leq \lambda \left(g(x) A \left(\gamma^{\frac{1}{p-1}} \frac{p-1}{p} \phi_{1,p}^{\frac{p}{p-1}} \right) + f \left(\gamma^{\frac{1}{q-1}} \frac{q-1}{q} \phi_{1,q}^{\frac{q}{q-1}} \right) \right), \end{aligned} \right.$$

then

$$\gamma r \left(\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right) \leq \lambda |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)].$$

By using (3.1)

$$\int_{\bar{\Omega}_\delta} |x|^{-ap} |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \omega_1 dx \leq \lambda \int_{\bar{\Omega}_\delta} |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)] \omega_1 dx. \quad (3.3)$$

Similarly, we shows that

$$\int_{\bar{\Omega}_\delta} |x|^{-bq} |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \omega_2 dx \leq \lambda \int_{\bar{\Omega}_\delta} |x|^{-(b+1)q+c_2} [g(x) B(\psi_2) + h(\psi_1)] \omega_2 dx. \quad (3.4)$$

Next, on $\Omega \setminus \bar{\Omega}_\delta$. Since $\lambda \geq A$, then

$$\lambda \geq \frac{\gamma \lambda_{1,p}}{\eta A \left(\gamma^{\frac{1}{p-1}} \alpha_p \right) + f \left(\gamma^{\frac{1}{q-1}} \alpha_q \right)},$$

thus, we have

$$\begin{aligned} \gamma \left(\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right) &\leq \gamma \lambda_{1,p} \\ &\leq \lambda \left[\eta A \left(\gamma^{\frac{1}{p-1}} \alpha_p \right) + f \left(\gamma^{\frac{1}{q-1}} \alpha_q \right) \right] \end{aligned}$$

and

$$\gamma r \left(\lambda_{1,p} |x|^{-(a+1)p+c_1} \phi_{1,p}^p - |x|^{-ap} |\nabla \phi_{1,p}|^p \right) \leq \lambda |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)], \quad (3.5)$$

so by (3.1)

$$\int_{\Omega \setminus \bar{\Omega}_\delta} |x|^{-ap} |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \omega_1 dx \leq \lambda \int_{\Omega \setminus \bar{\Omega}_\delta} |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)] \omega_1 dx. \quad (3.6)$$

Similarly, we shows that for all $\lambda \in [A, B]$

$$\int_{\Omega \setminus \bar{\Omega}_\delta} |x|^{-bq} |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \omega_2 dx \leq \lambda \int_{\Omega \setminus \bar{\Omega}_\delta} |x|^{-(b+1)q+c_2} [g(x) B(\psi_2) + h(\psi_1)] \omega_2 dx \quad (3.7)$$

(3.3) and (3.6) give:

$$\int_{\Omega} |x|^{-ap} |\nabla \psi_1|^{p-2} \nabla \psi_1 \nabla \omega_1 dx \leq \lambda \int_{\Omega} |x|^{-(a+1)p+c_1} [g(x) A(\psi_1) + f(\psi_2)] \omega_1 dx. \quad (3.8)$$

Similarly, we shows that

$$\int_{\Omega} |x|^{-bq} |\nabla \psi_2|^{q-2} \nabla \psi_2 \nabla \omega_2 dx \leq \lambda \int_{\Omega} |x|^{-(b+1)q+c_2} [g(x) B(\psi_2) + h(\psi_1)] \omega_2 dx. \quad (3.9)$$

From (3.8) and (3.9), we deduce that (ψ_1, ψ_2) is a sub-solution of (1.1). However, we have $\psi_1 > 0$ and $\psi_2 > 0$ in Ω .

Next, we introduce a supersolution of problem (1.1). For this intent, we can prove that there exists a large enough positive constant C so that

$$(z_1, z_2) = \left(\frac{C}{\mu_p} \lambda^{\frac{1}{p-1}} \zeta_p, \left[2h \left(C \lambda^{\frac{1}{q-1}} \right) \right]^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \zeta_q \right).$$

Let $\omega_1 \in W_0^{1,p}(\Omega, |x|^{-ap})$ with $\omega_1 \geq 0$.

For sufficient C large

$$\frac{\mu_p^{p-1} \left[\|g\|_\infty A \left(C \lambda^{\frac{1}{p-1}} \right) + f \left(2h \left(C \lambda^{\frac{1}{q-1}} \right) \right)^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \mu_q \right]}{C^{p-1}} \leq 1,$$

then

$$\begin{aligned} \int_{\Omega} |x|^{-ap} |\nabla z_1|^{p-2} \nabla z_1 \nabla \omega_1 dx &= \lambda \left(\frac{C}{\mu_p} \right)^{p-1} \int_{\Omega} |x|^{-ap} |\nabla \zeta_p|^{p-2} \nabla \zeta_p \nabla \omega_1 dx \\ &= \lambda \left(\frac{C}{\mu_p} \right)^{p-1} \int_{\Omega} |x|^{-(a+1)p+c_1} \omega_1 dx \\ &\geq \lambda \int_{\Omega} |x|^{-(a+1)p+c_1} \left[\|g\|_\infty A \left(C \lambda^{\frac{1}{p-1}} \right) + f \left(2h \left(C \lambda^{\frac{1}{q-1}} \right) \right)^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \mu_q \right] \omega_1 dx \quad (3.10) \\ &\geq \lambda \int_{\Omega} |x|^{-(a+1)p+c_1} \left[\|g\|_\infty A \left(C \lambda^{\frac{1}{p-1}} \frac{\zeta_p}{\mu_p} \right) + f \left(2h \left(C \lambda^{\frac{1}{q-1}} \right) \right)^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \zeta_q \right] \omega_1 dx \\ &= \lambda \int_{\Omega} |x|^{-(a+1)p+c_1} [g(x) A(z_1) + f(z_2)] \omega_1 dx. \end{aligned}$$

Similarly, we choose C large so that

$$\frac{\|g\|_\infty \left(B \left(2h \left(C \lambda^{\frac{1}{p-1}} \right) \right)^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \mu_q \right)}{h \left(C \lambda^{\frac{1}{p-1}} \right)} \leq 1.$$

Let $\omega_1 \in W_0^{1,p}(\Omega, |x|^{-ap})$ with $\omega_1 \geq 0$, then

$$\begin{aligned}
 \int_{\Omega} |x|^{-bq} |\nabla z_2|^{q-2} \nabla z_2 \nabla \omega_2 dx &= 2\lambda h \left(C \lambda^{\frac{1}{p-1}} \right) \int_{\Omega} |x|^{-bq} |\nabla \zeta_q|^{p-2} \nabla \zeta_q \nabla \omega_2 dx \\
 &= 2\lambda h \left(C \lambda^{\frac{1}{p-1}} \right) \int_{\Omega} |x|^{-(b+1)q+c_2} \omega_2 dx \\
 &\geq \lambda \int_{\Omega} |x|^{-(b+1)q+c_2} \|g\|_{\infty} \left(B \left(2h \left(C \lambda^{\frac{1}{p-1}} \right) \right)^{\frac{1}{q-1}} \lambda^{\frac{1}{q-1}} \mu_q \right) \omega_2 dx \\
 &= \lambda \int_{\Omega} |x|^{-(b+1)q+c_2} [g(x) B(z_2) + h(z_1)] \omega_2 dx.
 \end{aligned} \tag{3.11}$$

From (3.10) and (3.11) yield that (z_1, z_2) is a super-solution of problem (1.1) with $\psi_1 \leq z_1$ and $\psi_2 \leq z_2$ for $C > 0$ large.

Hence by Lemma (1.1), there exist a positive solution (u, v) of (1.1), where

$$(\psi_1, \psi_2) \leq (u, v) \leq (z_1, z_2).$$

ACKNOWLEDGEMENT

The authors would like to thank the anonymous referees and the handling editor for their careful reading. The first author gratefully acknowledge Qassim University in Kingdom of Saudi Arabia and this presented work is in memory of his father (1910-1999) Mr. Mahmoud ben Mouha Boulaaras.

REFERENCES

1. G. A. Afrouzi and S. H. Rasouli, A remark on the existence of multiple solutions to a multiparameter nonlinear elliptic system, *Nonlinear Anal.*, **71** (2009), 445-455.
2. G. A. Afrouzi and S. Haghaieghi, Sub-super solutions for $(p - q)$ Laplacian systems, *Boundary Value Problems*, 2011 2011:52.
3. G. A. Afrouzi, D., Vicențiu, E. Rădulescu, and S. Shaker, *i*Positive solutions of singular elliptic systems with multiple parameters and Caffarelli–Kohn–Nirenberg exponents, **70**(2) (2015), 145-152.
4. N. Akhmediev and A. Ankiewicz, Partially coherent solitons on a finite background, *Phys. Rev. Lett.*, **82** (1999), 2661-2664.
5. J. Ali and R. Shivaji, Positive solutions for a class of p -Laplacian systems with multiple parameters, *J. Math. Anal. Appl.*, **335** (2007), 1013-1019.

6. J. Ali, R. Shivaji, and M. Ramaswamy, Multiple positive solutions for classes of elliptic systems with combined nonlinear effects, *Differ. Integral Equ.*, **19** (2006), 669-680.
7. C. O. Alves and D. G. de Figueiredo, Nonvariational elliptic systems, *Discrete Contin. Dyn. Syst.*, **8** (2002), 289-302.
8. A. Ambrosetti, J. G. Azorero, and I. Peral, Existence and multiplicity results for some nonlinear elliptic equations, *Rend. Mat. Appl.*, **7** (2000), 167-198.
9. C. Atkinson and K. El Kalli, Some boundary value problems for the Bingham model, *J. Non Newton. Fluid Mech.*, **41** (1992), 339-363.
10. S. Boulaaras, R. Ghfaifia, and S. Kabli, An asymptotic behavior of positive solutions for a new class of elliptic systems involving of $(p(x), q(x))$ -Laplacian systems, *Bol. Soc. Mat. Mex.*, (2017).
<https://doi.org/10.1007/s40590-017-0184-4>.
11. H. Bueno, G. Ercole, W. Ferreira, and A. Zumpano, Existence and multiplicity of positive solutions for the p -Laplacian with nonlocal coefficient, *J. Math. Anal. Appl.*, **343** (2008), 151-158.
12. L. Caffarelli, R. Kohn, and L. Nirenberg, First order interpolation inequalities with weights, *Compos. Math.*, **53** (1984), 259-275.
13. A. Canada, P. Drábek, and J. L. Gámez, Existence of positive solutions for some problems with nonlinear diffusion, *Trans. Am. Math. Soc.*, **349** (1997), 4231-4249.
14. F. Catrina and Z.-Q. Wang, On the Caffarelli-Kohn-Nirenberg inequalities: Sharp constants, existence (and nonexistence), and symmetry of extremal functions, *Commun. Pure Appl. Math.*, **54**(2001), 229-258.
15. R. Dalmaso, Existence and uniqueness of positive solutions of semilinear elliptic systems, *Nonlinear Anal.*, **39** (2000), 559-568.
16. E. N. Dancer, Competing species systems with diffusion and large interaction, *Rend. Sem. Mat. Fis. Milano*, **65** (1995), 23-33.
17. R. Dautray and J.-L. Lions, *Mathematical analysis and numerical methods for science and technology*, **1**: Physical Origins and Classical Methods, Springer, Berlin, Heidelberg, NewYork, 1985.
18. P. Drabek and J. Hernandez, Existence and uniqueness of positive solutions for some quasilinear elliptic problem, *Nonlinear Anal.*, **44** (2001), 189-204.
19. J. F. Escobar, Uniqueness theorems on conformal deformations of metrics, Sobolev inequalities, and an eigenvalue estimate, *Commun. Pure Appl. Math.*, **43** (1990), 857-883.
20. F. Fang and S. Liu, Nontrivial solutions of superlinear p -Laplacian equations, *J. Math. Anal. Appl.*, **351** (2009), 3601-3619.
21. R. Filippucci, P. Pucci, and V. Radulescu, Existence and non-existence results for quasilinear elliptic exterior problems with nonlinear boundary conditions, *Commun. Partial Differ. Equ.*, **33** (2008), 706-717.

22. D. D. Hai and R. Shivaji, An existence result on positive solutions for a class of semi-linear elliptic systems, *Proc. R. Soc. Edinb. Sect. A*, **134** (2004), 137-141.
23. R. Ghfaifia and S. Boulaaras, Existence of positive solution for a class of $(p(x); q(x))$ -Laplacian systems, *Rend. Circ. Mat. Palermo, II. Ser.*, **67** (2018), 93-103. <https://doi.org/10.1007/s12215-017-0297-7>.
24. J. R. Graef, S. Heidarkhani, and L. Kong, Multiple solutions for systems of multi-point boundary value problems, *Opusc. Math.*, **33** (2013), 293-306.
25. D. D. Hai and R. Shivaji, An existence result on positive solutions for a class of p -Laplacian systems, *Nonlinear Anal.*, **56** (2004), 1007-1010.
26. O. H. Miyagaki and R. S. Rodrigues, On positive solutions for a class of singular quasilinear elliptic systems, *J. Math. Anal. Appl.*, **334** (2007), 818-833.
27. M. K. V. Murthy and G. Stampacchia, Boundary value problems for some degenerate elliptic operators, *Ann. Mat. Pura Appl.*, **80** (1968), 1-122.
28. M. Nagumo, Über die Differentialgleichung $y = f(x, y, y)$, *Proc. Phys. Math. Soc. Jpn.*, **19** (1937), 861-866.
29. H. Poincaré, Les fonctions fuchsiennes et l'équation $u = e^u$, *J. Math. Pures Appl.*, **4** (1898), 137-230.
30. P. Tolksdorf, Regularity for a more general class of quasilinear elliptic equations, *J. Diff. Equ.*, **51** (1984), 126-150.
31. B. Xuan, The eigenvalue problem for a singular quasilinear elliptic equation, *Electron. J. Differ. Equ.*, **16** (2004), 1-11.