

NUMERICAL IMPLEMENTATION OF FINITE-TIME SHADOWING OF STOCHASTIC DIFFERENTIAL EQUATIONS

Qingyi Zhan^{*,**}, Zhifang Zhang^{***} and Yuhong Li^{****}

**College of Computer and Information Science, Fujian Agriculture and Forestry University,
Fuzhou, 350002, P.R. China*

***Department of Applied Mathematics, Illinois Institute of Technology,
Chicago, IL, 60616, USA*

****Fujian Center for Disease Control and Prevention, Fuzhou,
Fujian, 350001, P.R. China*

*****College of Hydropower and Information Engineering, Huazhong University of
Science and Technology, Wuhan, 430074, P.R. China*

e-mails: zhanqy@lsec.cc.ac.cn, qzhan3@iit.edu; liyuhong@hust.edu.cn

*(Received 16 September 2018; after final revision 11 December 2019;
accepted 31 January 2020)*

This paper focuses on the numerical implementation methods of shadowing theorem of stochastic differential equations. A general shadowing theorem of stochastic differential equations is given, and an explicit bound for shadowing distance is investigated. The main part is numerical implementation methods for shadowing distance in details. Numerical experiments are provided to illustrate the effectiveness of the proposed theorem by the numerical simulations of chaotic orbits of stochastic differential equations.

Key words : Stochastic differential equations, random shadowing, numerical implementation, stochastic Chen system.

2010 Mathematics Subject Classification : 37H05, 37C50, 65C30, 65P20.

1. INTRODUCTION

Nowadays, the shadowing property has played an important role in the theory and application of random dynamical systems (RDS), especially in numerical simulations of chaotic systems of stochastic differential equations (SDEs). Numerical computation is in the center in the investigations of chaotic

SDEs which describe many natural phenomena, for example, in meteorology and biology [1, 3, 11, 12, 17]. However, we mainly concern the reliability and feasibility of numerical computations. These related works are shown in References [8, 19-22].

This work is motivated by two facts. Firstly, the construction of conditions which can assure stochastic shadowing in a class of SDEs has been finished in [19]. Many useful contributions are made to the numerical analysis of RDS [14, 15, 16]. These results are the foundations of stochastic shadowing. Secondly, the high accurate numerical implementation methods of stochastic shadowing continues to be an interested topic. To the best of our knowledge, no investigations of the numerical implementations of random shadowing of SDEs in a finite time interval exist in the literature.

In this work, we mainly focus on the numerical methods to compute all parameters which are involved in a shadowing theorem of SDEs in a finite time interval. This is the continuation of Paper [19-22] in some sense. The results show that under certain appropriate assumptions the numerical approximate orbits of discrete approximate RDS are more close to the true orbits of the original systems.

The rest of this paper is organized as follows. Section 2 deals with some preliminaries. In Section 3 the theoretical results of the finite time shadowing are summarized. Section 4 presents the details of the numerical implementations. Illustrative numerical experiments for the main theorem are included in Section 5. We demonstrate that the main theorem and numerical implementation methods can be applied to stochastic Chen system (SCS), stochastic Lorenz equations (SLEs) and stochastic Lü equations (SLuEs). Finally, section 6 is addressed to summarize the conclusions of the paper.

2. PRELIMINARIES

Let $W_t, t \in \mathbb{R}^+ := [0, +\infty)$ be a one dimensional Wiener process, and $(\Omega, \mathcal{F}, \mathbb{P})$ be a Wiener space. And we assume that $\Omega := \{\omega \in C(\mathbb{R}^+, \mathbb{R}) : \omega(0) = 0\}$. We consider a class of Stratonovich SDEs as follows,

$$dx_t = f_0(x_t)dt + f_1(x_t) \circ dW_t, \quad x(0) = x_0(\omega) \in \mathbb{R}^d, \quad (1)$$

where the random variable $x_0(\omega)$ is independent of $\mathcal{F}_0$ and satisfies the inequality $\mathbb{E}|x_0(\omega)|^2 < \infty$. And $f_0 \in \mathcal{C}_b^{1, \delta_1}, f_1 \in \mathcal{C}_b^{2, \delta_1}, \sum_{i=1}^d f_1^i \frac{\partial f_1}{\partial x_i} \in \mathcal{C}_b^{1, \delta_1}$, where the space $\mathcal{C}_b^{k, \delta_1}$ is defined to be the Banach space of function $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$, which are in $\mathcal{C}^{k, \delta_1}$ and for which the norm

$$\|f\|_{k,0} := \sup_{x \in \mathbb{R}^d} \frac{\|f(x)\|}{1 + \|x\|} + \sum_{1 \leq |\alpha| \leq k} \sup_{x \in \mathbb{R}^d} |D^\alpha f(x)|,$$

and

$$\|f\|_{k,\delta_1} := \|f\|_{k,0} + \sum_{|\alpha|=k} \sup_{x \neq y} \frac{\|D^\alpha f(x) - D^\alpha f(y)\|}{\|x - y\|^{\delta_1}}, 0 < \delta_1 \leq 1, k = 1, 2,$$

is finite, see References [1, 4].

We define

$$\theta : \mathbb{R}^+ \times \Omega \rightarrow \Omega, \theta^t \omega(s) = \omega(t + s) - \omega(t)$$

and $0 \leq s \leq t$, $s \in \mathbb{R}^+$, $t \in \mathbb{R}^+$. It follows from Theorem 2.3.32 in [1] that SDE (1) generates a unique RDS $\varphi : \mathbb{R}^+ \times \mathbb{R}^+ \times \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^d$, which is usually written as $\varphi(s, t, \omega)x := \varphi(s, t, \omega, x)$ on the metric dynamical systems $(\Omega, \mathcal{F}, \mathbb{P}, \theta^\approx)$, and is C^1 with respect to x . The RDS φ is written as follows,

$$\varphi(s, t, \omega)x = x + \int_s^t f_0(\varphi(s, r, \omega)x)dr + \int_s^t f_1(\varphi(s, r, \omega)x) \circ dW_r(\omega).$$

We also make use of the following notations, the norm of a random variable $\|\cdot\|$, the norm of random matrix $\|\cdot\|$ and the norm of a stochastic process $\|x(t, \omega)\|_2$, which are similar to [18, 19].

3. THEORETICAL FOUNDATIONS OF NUMERICAL SHADOWING

3.1 Some existed definitions

Definition 3.1 — [19]. Let $\mathcal{F}_{t_k}$ be the filtration generated by $W_\tau(\omega)$, $0 \leq \tau \leq t_k$, and random variable $x_0(\omega)$ be independent of $\mathcal{F}_{t_0}$. For a given positive number δ and $\mathbb{P}$ -almost surely $\omega \in \Omega$, if there is a sequence of time $\{t_k\}_{k=0}^N$, $0 \leq t_0 \leq t_1 \leq \dots \leq t_N$ and a sequence of random variables $\{(u_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ which means that $u_k(\theta^{t_k}\omega)$ is $\mathcal{F}_{t_k}$ -measurable for $k = 0, 1, \dots, N$, such that the following inequalities hold

$$\|u_{k+1}(\theta^{t_{k+1}}\omega) - \varphi(t_k, t_{k+1}, \theta^{t_k}\omega)u_k(\theta^{t_k}\omega)\| \leq \delta, \quad k = 0, 1, \dots, N-1,$$

then the random variables $\{(u_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ are said to be a (ω, δ) -pseudo orbit of SDE (1) in the mean-square sense, where $\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)u_k(\theta^{t_k}\omega)$ denotes the orbit of RDS φ at the time t_{k+1} which starts from the initial time t_k with the initial value $u_k(\theta^{t_k}\omega)$ and the sample $\theta^{t_k}\omega$, and $u_k(\theta^{t_k}\omega)$ denotes a random variable with the sample $\theta^{t_k}\omega$ which is $\mathcal{F}_{t_k}$ -measurable.

Definition 3.2 — [19]. For a given positive number ε , $\mathbb{P}$ -almost surely $\omega \in \Omega$, and a (ω, δ) -pseudo orbit $\{(u_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ of SDE (1) with associated times $\{t_k\}_{k=0}^N$, if there is a sequence of time $\{h_k\}_{k=0}^N$, $0 \leq h_0 = t_0 \leq h_1 \leq \dots \leq h_N$, which is random, such that the following inequalities hold

$$\|u_k(\theta^{t_k}\omega) - x_k(\theta^{h_k}\omega)\| \leq \varepsilon, \quad 0 \leq t_k - h_k \leq \varepsilon, \quad k = 0, 1, \dots, N, \quad (2)$$

where the random variables $\{(x_k(\theta^{h_k}\omega), \mathcal{F}_{h_k})\}_{k=0}^N$ are on the true orbit of SDE (1), that is

$$x_{k+1}(\theta^{h_{k+1}}\omega) = \varphi(h_k, h_{k+1}, \theta^{h_k}\omega)x_k(\theta^{h_k}\omega), \quad k = 0, 1, \dots, N-1,$$

then the (ω, δ) -pseudo orbit $\{(u_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ is said to be (ω, ε) -shadowed by a true orbit of SDE (1) in the mean-square sense, where the true orbit of RDS φ is a stochastic process.

3.2 Theoretical results of finite time shadowing

Here we repeat some results in Reference [20]. For a (ω, δ) -pseudo orbit $\{(y_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$, we want to prove that it is (ω, ε) -shadowed by a true orbit containing points $\{(\hat{x}_k(\theta^{\hat{h}_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ for $\mathbb{P}$ -almost surely $\omega \in \Omega$, where $\{\hat{h}_k\}_{k=0}^N = \{t_k\}_{k=0}^N$ in sequels. For simplicity, we define that $t_k := k\Delta t$, Δt is the given time step, $\hat{x}_k(\theta^{\hat{h}_k}\omega)$ is the point on the true orbit, $\mathcal{F}_{t_k}$ is the filtration generated by $W_\tau(\omega)$, $0 \leq \tau \leq t_k$, and $N := \frac{t}{\Delta t}$.

We first introduce the function space $\mathcal{X}$ and $\mathcal{Y}$, which are the same as Reference [20]. Specially,

$$y(\omega) =: (y_0(\theta^{t_0}\omega), y_1(\theta^{t_1}\omega), \dots, y_N(\theta^{t_N}\omega)) \in \mathbb{R}^{(N+1)d},$$

is a special element of the space $\mathcal{Y}$. And

$$x(\omega) = (x_0(\theta^{h_0}\omega), x_1(\theta^{h_1}\omega), \dots, x_N(\theta^{h_N}\omega)) \in \mathbb{R}^{(N+1)d}$$

is a general element of the space $\mathcal{X}$.

Secondly, we define the mapping $G : \mathcal{X} \longrightarrow \mathcal{Y}$, and the k th component of $G(y(\omega))$ is defined for any $y(\omega) \in \mathcal{X}$ to be

$$[G(y(\omega))]_k = y_k(\theta^{t_k}\omega) - \varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega), \quad k = 1, \dots, N. \quad (3)$$

Here we consider the first variation $DG(y(\omega))$ of $G(y(\omega))$. By the assumptions that RDS φ generated by SDE (1) is C^1 with respect to x , then the Jacobian matrix $D\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}$ with respect to y_{k-1} exists. It follows from $G(y(\omega))$ is C^1 with respect to $y(\omega)$ that the existence of $DG(y(\omega))$ is obtained. Then $DG(y(\omega))$ is defined by the following equality

$$DG(y(\omega)) := \begin{pmatrix} -D\varphi(t_0, t_1, \theta^{t_0}\omega)y_0 & \mathbb{I} & 0 & \dots & 0 & 0 \\ 0 & -D\varphi(t_1, t_2, \theta^{t_1}\omega)y_1 & \mathbb{I} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \mathbb{I} & 0 \\ 0 & 0 & 0 & \dots & -D\varphi(t_{N-1}, t_N, \theta^{t_{N-1}}\omega)y_{N-1} & \mathbb{I} \end{pmatrix}, \quad (4)$$

where

$$D\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega) = \frac{\partial}{\partial y}(\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega)), \quad k = 1, 2, \dots, N,$$

that is, $DG(y(\omega))$ is a $Nd \times (N+1)d$ random matrix, and $\mathbb{I}$ is the $d \times d$ identity matrix.

Lastly, we wish to find a solution

$$\hat{x}(\omega) := (\hat{x}_0(\theta^{t_0}\omega), \hat{x}_1(\theta^{t_1}\omega), \dots, \hat{x}_N(\theta^{t_N}\omega)) \in \mathcal{X}, 0 \leq t_k - \hat{h}_k \leq \varepsilon$$

of the equation $G(\hat{x}(\omega)) = 0$ and to prove that it lies in a closed ε -neighborhood of $y(\omega)$.

Here we consider the first variation $DG(y(\omega))$ of $G(y(\omega))$. We define that

$$D\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega) = \frac{\partial}{\partial y}(\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega)), \quad k = 1, 2, \dots, N.$$

Then we have

$$[DG(y(\omega))\Delta y(\omega)]_k = \Delta y_k(\theta^{t_k}\omega) - D\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega) \cdot \Delta y_{k-1}(\theta^{t_{k-1}}\omega), \quad k = 1, 2, \dots, N,$$

where

$$\Delta y(\omega) = (\Delta y_0(\theta^{t_0}\omega), \Delta y_1(\theta^{t_1}\omega), \dots, \Delta y_N(\theta^{t_N}\omega))^T,$$

and

$$\Delta y_k(\theta^{t_k}\omega) = x_k(\theta^{t_k}\omega) - y_k(\theta^{t_k}\omega), \quad k = 0, 1, 2, \dots, N.$$

We assume that $B_k(\theta^{t_k}\omega)$ is a $d \times d$ random matrix obtained by numerical integration and satisfies

$$\|B_k(\theta^{t_k}\omega) - D\varphi(t_{k-1}, t_k, \theta^{t_{k-1}}\omega)y_{k-1}(\theta^{t_{k-1}}\omega)\| \leq \delta. \quad (5)$$

The detail of the numerical implementation will be shown in Section 4.1. Then we can obtain the numerical approximation $L(y(\omega))$ of $DG(y(\omega))$ and

$$[L(y(\omega))\Delta y(\omega)]_k = \Delta y_k(\theta^{t_k}\omega) - B_k(\theta^{t_k}\omega)\Delta y_{k-1}(\theta^{t_{k-1}}\omega), \quad k = 1, 2, \dots, N. \quad (6)$$

Theorem 3.3 — Let $G : \mathcal{X} \rightarrow \mathcal{Y}$ be a C^1 map as (3) and $L(y(\omega))$ be the numerical approximation of $DG(y(\omega))$, respectively. And let $y(\omega) \in \mathcal{X}$ be a point such that the pseudo inverses $[DG(y(\omega))]^{-1}$ and $[L(y(\omega))]^{-1}$ exist, and the following inequalities hold

$$\|G(y(\omega))\| \leq \delta,$$

$$\|[L(y(\omega))]^{-1}\| \leq c, \quad (7)$$

and

$$\|[L(y(\omega))]^{-1} - [DG(y(\omega))]^{-1}\| \leq \alpha \quad (8)$$

for some positive constants δ, c and α .

Furthermore, we let $\varepsilon := 2(\alpha + c)\delta$. If the inequality

$$\|DG(\hat{x}(\omega)) - L(y(\omega))\| = \beta \leq \frac{1}{2(\alpha + c)}, \quad (9)$$

holds for $\|\hat{x}(\omega) - y(\omega)\| \leq \varepsilon$, then the equation $G(\hat{x}(\omega)) = 0$ has one solution $\hat{x}(\omega)$ which satisfies $\|\hat{x}(\omega) - y(\omega)\| \leq \varepsilon$.

PROOF : The proof is similar to Reference [20], so we omit the detail here. $\square$

4. NUMERICAL IMPLEMENTATION METHODS

This section is the main part of this article. We approximate the local error δ using the local error control mechanism of the numerical scheme. Then we only pay attention to the magnification of the local error, $(\alpha + c)$, that gives the shadowing distance. This is part quote from Paper [20].

4.1 Basic method

Step 1 : Applying one-step numerical schemes (EM scheme or Milstein scheme [9]) to simultaneously solve the following equations from t_k to t_{k+1} with the initial values $x(0) = y_k(\theta^{t_k}\omega)$ and $v(0) = \mathbb{I}$,

$$\begin{cases} dx_t = f_0(x_t)dt + f_1(x_t) \circ dW_t(\omega) \\ dv_t = Df_0(x_t)v_t dt + Df_1(x_t)v_t \circ dW_t(\omega), \end{cases}$$

we get the approximations of $\hat{x}_{k+1}(\theta^{t_{k+1}}\omega)$ and $D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)$ respectively,

$$\hat{x}_{k+1}(\theta^{t_{k+1}}\omega) \approx \varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega),$$

$$D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) \approx v_{k+1}(\theta^{t_{k+1}}\omega) = B_{k+1}(\theta^{t_{k+1}}\omega).$$

That is, $B_{k+1}(\theta^{t_{k+1}}\omega) := D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)$ is chosen such that the inequality (5) hold by numerical schemes.

Step 2 : It follows from Theorem 3.3 that $[L(y(\omega))]^{-1}$ and $[DG(y(\omega))]^{-1}$ both exist. By the oscillation theorem of linear equations, we can find α such that

$$\|[L(y(\omega))]^{-1} - [DG(y(\omega))]^{-1}\| \leq \alpha.$$

The detail of this estimate will be shown in Section 4.2. And the estimate of the parameter c will be shown in Section 4.2, too.

Step 3 : By the norm properties, we can find β such that

$$\beta = \|DG(\hat{x}(\omega)) - L(y(\omega))\|.$$

If the inequality (3.1) is not satisfied, then we will denote this by putting '—' in the table of numerical experiment results. And the detail of the method will be shown in Section 4.3.

Step 4 : If the values of c, α, δ and β satisfy all conditions of Theorem 3.1, then the shadowing distance is $\varepsilon = 2(c + \alpha)\delta$.

4.2 Methods to estimate the parameters α and c

Let $A := L(y(\omega))[L(y(\omega))]^T$ and $A + \Delta A := DG(y(\omega))[DG(y(\omega))]^T$. Since $[L(y(\omega))]^{-1} = [L(y(\omega))]^T A^{-1}$, then $[L(y(\omega))]^{-1}$ and $[DG(y(\omega))]^{-1}$ can be formed by the solutions of the following equations, respectively,

$$Ax = b^{(i)}, \quad (A + \Delta A)y = b^{(i)} + \Delta b^{(i)},$$

where $\Delta A = DG(y(\omega))[DG(y(\omega))]^T - L(y(\omega))[L(y(\omega))]^T$, and $b^{(i)} = [L(y(\omega))]_i$ is the i -th column vector of $L(y(\omega))$, $i = 1, 2, \dots, (N+1)d$. Furthermore, $\Delta b^{(i)} := \Delta[L(y(\omega))]_i = [L(y(\omega))]_{i+1} - [L(y(\omega))]_i$.

The following lemma is a direct consequence in [6].

Lemma 4.1 — Suppose $Ax = b^{(i)}$, $A \in \mathbb{R}^{Nd \times Nd}$, $0 \neq b^{(i)} \in \mathbb{R}^{Nd \times 1}$, and $(A + \Delta A)y = b^{(i)} + \Delta b^{(i)}$, $\Delta A \in \mathbb{R}^{Nd \times Nd}$, $\Delta b^{(i)} \in \mathbb{R}^{Nd \times 1}$, with $\|\Delta A\|_1 \leq \zeta \|A\|_1$, $\|\Delta b^{(i)}\|_1 \leq \zeta \|b^{(i)}\|_1$. If $\zeta \kappa(A) < 1$, then $A + \Delta A$ is nonsingular and

$$\frac{\|y - x\|_1}{\|x\|_1} \leq \frac{2\zeta}{1 - \zeta \kappa(A)} \kappa(A),$$

where $\kappa(A)$ is the condition number of the matrix A , $\zeta > 0$, $\|\cdot\|_1$ is the 1-norm.

Proposition 4.2 — Suppose $\{(y_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ is a (ω, δ) -pseudo orbit of SDE (1). If ζ and $\kappa(A)$ satisfy the conditions of Lemma 4.1, then the following conclusions hold

$$\|[L(y(\omega))]^{-1}\| \leq c \text{ and } \alpha = \frac{2\zeta}{1 - \zeta \kappa(A)} \kappa(A)c,$$

where

$$\zeta = \max\{\zeta_1, \zeta_2\}$$

and

$$\zeta_1 = \sup_i \left\{ \frac{\delta}{\|b^{(i)}\|_1} \right\}, \zeta_2 = \sup_k \left\{ \frac{\delta(\|B_k(\theta^{t_k}\omega)\|_1 + \|B_k(\theta^{t_k}\omega)\|_\infty + \delta + 2)}{\|A\|_1} \right\}. \quad (10)$$

PROOF : Firstly, it follows from Section 3.2 that $DG(y(\omega))$ exists. Furthermore, Theorem 3.3 shows the existence of $[L(y(\omega))]^{-1}$. Therefore, by the definition of the norm the constant c does exist. And the estimate of $\|[L(y(\omega))]^{-1}\|$ is similar to Reference [20].

Secondly, we estimate the parameter α [20]. On the one hand, note that the quantities $\{(y_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$ and $L(y(\omega))$ can be calculated by one-step numerical scheme as shown in Section 4.1. Then the condition number $\kappa(A)$ can be found by the Matlab code during our computations. We need to estimate the supremum norm of $L(y(\omega))$. It follows from (6) and the definition of the 1-norm that $\|\Delta b^{(i)}\|_1 := \|B_{i+1}(\theta^{t_{i+1}}\omega) - B_i(\theta^{t_i}\omega)\|_1 = \delta$. Note that we can calculate every row of $b^{(i)}$ and obtain $\zeta_1 = \max \left\{ \frac{\delta}{\|b^{(i)}\|_1} \right\}$ such that $\|\Delta b^{(i)}\|_1 \leq \zeta_1 \|b^{(i)}\|_1$, where

$$\|b^{(i)}\|_1 = \sum_{j=1}^{Nd} \|(L(y(\omega)))_{ij}\|.$$

On the other hand, we have

$$\|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k\|_1 \leq \|B_k(\theta^{t_k}\omega)\|_1 + \delta.$$

Therefore, it follows from the property of norm that we get

$$\begin{aligned} & \|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k \cdot [D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k]^T - B_k(\theta^{t_k}\omega) \cdot [B_k(\theta^{t_k}\omega)]^T\|_1 \\ &= \|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k \cdot [D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k]^T \\ & - [B_k(\theta^{t_k}\omega) - D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k + D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k] \cdot [B_k(\theta^{t_k}\omega)]^T\|_1 \\ &\leq \|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k \cdot [D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k]^T - [B_k(\theta^{t_k}\omega)]^T\|_1 \\ &+ \|[D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k - B_k(\theta^{t_k}\omega)] \cdot [B_k(\theta^{t_k}\omega)]^T\|_1 \\ &\leq \delta(\|B_k(\theta^{t_k}\omega)\|_1 + \|B_k(\theta^{t_k}\omega)\|_\infty + \delta) := C_1^{(k)}, \end{aligned}$$

where $\|[B_k(\theta^{t_k}\omega)]^T\|_1 \leq \|B_k(\theta^{t_k}\omega)\|_\infty$, and $\|\cdot\|_\infty$ is the infinite norm.

Note that A is block tridiagonal, it follows from the random matrix computation that we obtain

$$\|\Delta A\|_1 = \sup_k \{2\delta + C_1^{(k)}\}$$

and the second part of (10) such that $\|\Delta A\|_1 \leq \zeta_2 \|A\|_1$, where

$$\|B_k(\theta^{t_k}\omega)\|_\infty = \max \left\{ \sum_{j=1}^d \|(B_k(\theta^{t_k}\omega))_{1j}\|, \sum_{j=1}^d \|(B_k(\theta^{t_k}\omega))_{2j}\|, \dots, \sum_{j=1}^d \|(B_k(\theta^{t_k}\omega))_{dj}\| \right\}.$$

We choose $\zeta = \max\{\zeta_1, \zeta_2\}$ such that $\|\Delta b^{(i)}\|_1 \leq \zeta \|b^{(i)}\|_1$ and $\|\Delta A\|_1 \leq \zeta \|A\|_1$. It follows from Lemma 4.1 that

$$\alpha = \frac{2\zeta}{1 - \zeta\kappa(A)} \kappa(A)c.$$

This proof is finished. $\square$

4.3 — Method to estimate the parameter β

This part will prove that the parameter β can be explicitly estimated.

Proposition 4.3 — Assume that the functions f_0 and f_1 satisfies the conditions in Section 2. In the proper ε_0 -neighbourhood of the (ω, δ) -pseudo orbit $\{(y_k(\theta^{t_k}\omega), \mathcal{F}_{t_k})\}_{k=0}^N$, let the local Lipschitz constants of f_0 , f_1 and f_2 be K_1 , and the local Lipschitz constants of Df_0 and Df_1 be K_2 , where the parameters K_1 and K_2 are positive and independent of δ and t_k , and $f_2 = f_0 + \frac{1}{2}f_1'f_1$.

Then the following equality holds

$$\beta = K_3 + \delta,$$

where

$$K_3 = \sup_k K_2 \sqrt{2\Delta t_{k+1}(\Delta t_{k+1} + 1)K_4}, \quad (11)$$

$$K_4 = \sup_k \left[3\varepsilon_0^2 \exp[3\Delta t_{k+1}(\Delta t_{k+1} + 1)K_1^2] \right] \text{ and } \Delta t_{k+1} = t_{k+1} - t_k. \quad (12)$$

PROOF : It follows from the definition of DG that

$$\|DG(\hat{x}(\omega)) - L(y(\omega))\| \leq K_5 + \delta,$$

where

$$K_5 = \sup_k \|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega) - D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)\|.$$

First of all, we estimate the error caused by different initial value. The constant $\varepsilon_0 \in (0, \varepsilon]$ is given such that the inequality $\|y_k(\theta^{t_k}\omega) - \hat{x}_k(\theta^{t_k}\omega)\| \leq \varepsilon_0$ hold, then we have

$$\mathbb{E} \left[\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega) \right]^2 \leq I_1 + I_2 + I_3,$$

where

$$I_1 = 3\mathbb{E}\left[y_k(\theta^{t_k}\omega) - \hat{x}_k(\theta^{t_k}\omega)\right]^2,$$

$$I_2 = 3\mathbb{E}\left[\int_{t_k}^{t_{k+1}} [f_2(\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)) - f_2(\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega))]ds\right]^2$$

and

$$I_3 = 3\mathbb{E}\left[\int_{t_k}^{t_{k+1}} [f_1(\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)) - f_1(\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega))]dW_s(\omega)\right]^2.$$

Utilizing Cauchy-Schwarz inequality and the local Lipschitz condition of f_2 , we have

$$I_2 \leq 3\Delta t_{k+1}K_1^2 \int_{t_k}^{t_{k+1}} \mathbb{E}\left[\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 ds.$$

It follows from the local Lipschitz condition of f_1 and the Itô isometry that we have

$$I_3 \leq 3K_1^2 \int_{t_k}^{t_{k+1}} \mathbb{E}\left[\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 ds.$$

Therefore, we obtain

$$\begin{aligned} & \mathbb{E}\left[\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 \\ & \leq 3\mathbb{E}\left[y_k(\theta^{t_k}\omega) - \hat{x}_k(\theta^{t_k}\omega)\right]^2 \\ & + 3(\Delta t_{k+1} + 1)K_1^2 \int_{t_k}^{t_{k+1}} \mathbb{E}\left[\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 ds. \end{aligned}$$

It follows from the Gronwall inequality and (12) that we obtain

$$\mathbb{E}\left[\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 \leq K_4. \quad (13)$$

Secondly, we estimate the value of K_5 . The expression K_5 implies

$$\mathbb{E}\left[D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 \leq I_4 + I_5,$$

where

$$\begin{aligned} I_4 = 2\mathbb{E}\left[\int_{t_k}^{t_{k+1}} [Df_0(\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)) \cdot D\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) \right. \\ \left. - Df_0(\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)) \cdot D\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)]ds\right]^2 \end{aligned}$$

and

$$I_5 = 2\mathbb{E}\left[\int_{t_k}^{t_{k+1}} [Df_1(\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega)) \cdot D\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - Df_1(\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)) \cdot D\varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)] \circ dW_s(\omega)\right]^2.$$

Utilizing the Cauchy-Schwarz inequality and the local Lipschitz condition of Df_0 , we have

$$I_4 \leq 2\Delta t_{k+1} K_2^2 \int_{t_k}^{t_{k+1}} \mathbb{E}\left[\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 ds.$$

It follows from the Itô isometry and the local Lipschitz condition of Df_1 that we have

$$I_5 \leq 2K_2^2 \int_{t_k}^{t_{k+1}} \mathbb{E}\left[\varphi(t_k, s, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - \varphi(t_k, s, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 ds.$$

Using these conclusions and (13), we obtain

$$\mathbb{E}\left[D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\right]^2 \leq 2\Delta t_{k+1}(\Delta t_{k+1} + 1)K_2^2 K_4. \quad (14)$$

Therefore, by (11) and (14) we arrive at the inequality

$$\|D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)y_k(\theta^{t_k}\omega) - D\varphi(t_k, t_{k+1}, \theta^{t_k}\omega)\hat{x}_k(\theta^{t_k}\omega)\| \leq K_3. \quad (15)$$

By the conclusions of (11) and (15), we obtain $K_5 \leq K_3$. Therefore, we only need to choose

$$\beta = K_3 + \delta.$$

This completes the proof. $\square$

5. NUMERICAL EXPERIMENTS

5.1 Experimental preparation

We consider the Stratonovich SCS

$$dX_t = f_0(X_t)dt + f_1(X_t) \circ dW_t(\omega), \quad X(0) = x_0 \in \mathbb{R}^3 \quad (16)$$

where $X_t = (x(t), y(t), z(t))^T \in \mathbb{R}^3$, $x(t)$, $y(t)$ and $z(t)$ make up the system state, σ , ρ and β are the system parameters, and

$$f_0(X_t) = \begin{pmatrix} -\sigma x(t) + \sigma y(t) \\ (\rho - \sigma)x(t) + \rho y(t) - x(t)z(t) \\ -\beta z(t) + x(t)y(t) \end{pmatrix}, \quad f_1(X_t) = \begin{pmatrix} \mu x(t) \\ \mu y(t) \\ \mu z(t) \end{pmatrix}.$$

In this experiment, we choose the classic parameters $\sigma = 35, \rho = 28, \beta = 3$. A RDS (φ, θ) can be generated by the solution operator of SDE (16).

The preset initial conditions are as follows: the initial value is $x_0 = -10.0, y_0 = 1.0, z_0 = 37.0$. SDE (16) is discretized by Euler-Maruyama(EM) scheme, we obtain that

$$\begin{cases} x_{k+1} = x_k + (\sigma(-x_k + y_k) + 0.5\mu^2 x_k)\Delta t_k + \mu x_k \Delta W_k, \\ y_{k+1} = y_k + (-x_k z_k - (\rho - \sigma)x_k + \rho y_k + 0.5\mu^2 y_k)\Delta t_k + \mu y_k \Delta W_k, \\ z_{k+1} = z_k + (x_k y_k - \beta z_k + 0.5\mu^2 z_k)\Delta t_k + \mu z_k \Delta W_k. \end{cases} \quad (17)$$

5.2 Numerical results

This section will provide numerical experiments to compute the shadowing distance of SDE (16). The classical parameters are chosen and the initial value is $x_0 = -10.0, y_0 = 1.0, z_0 = 37.0$.

Firstly, in order to show the influence of noise on the pseudo orbits, we choose various size of noise, such as $\mu = 0$ (the deterministic case), $\mu = 1.0, \mu = 2.0$ and $\mu = 4.0$. And we take the temporal step-size $\Delta t = 1.05e - 3$ and $N = 1.0e + 4$. Taking the numerical solution for an example, Fig. 1 shows the perturbation of the pseudo orbits of SDE (16) by EM scheme corresponding to different scales of noise. As we can see from Fig. 1, due to the increase of the noise scale, that is, the value of μ is increased from 0 to 4.0, the perturbation of the pseudo orbits becomes much more seriously in x and y directions, respectively. That is to say, the pseudo orbits in x direction of Fig. 1 become more and more dense, which illustrates that the perturbation of the pseudo orbits in x direction become more and more larger. It is similar to the case in y direction.

Secondly, we focus on the numerically performing the shadowing distance, $\varepsilon = 2(\alpha + c)\delta$. The amplification factors α and c are computed quite accurately, and the local error tolerance δ is approximately determined by the numerical scheme. Applying EM scheme and Milstein scheme to SCS, we present the discrete forms respectively.

SDE (16) is discretized by Milstein scheme as follows,

$$\begin{cases} x_{k+1} = x_k + (\sigma(-x_k + y_k) + 0.5\mu^2 x_k)\Delta t_k + \mu x_k \Delta W_k + 0.5\mu^2 x_k((\Delta W_k)^2 - \Delta t_k), \\ y_{k+1} = y_k + (-x_k z_k - (\rho - \sigma)x_k + \rho y_k + 0.5\mu^2 y_k)\Delta t_k + \mu y_k \Delta W_k + 0.5\mu^2 y_k((\Delta W_k)^2 - \Delta t_k), \\ z_{k+1} = z_k + (x_k y_k - \beta z_k + 0.5\mu^2 z_k)\Delta t_k + \mu z_k \Delta W_k + 0.5\mu^2 z_k((\Delta W_k)^2 - \Delta t_k). \end{cases} \quad (18)$$

Table 1 presents the results that δ is the local error and ε is the shadowing distance. We can obtain that the existence of shadowing orbit and the effectiveness of the numerical method. redAs we know, the shadowing error δ obtained by Milstein scheme is smaller than that by EM scheme. Therefore,




Fig. 1. Pseudo orbits for different sizes of noise $\mu = 0, \mu = 1.0, \mu = 2.0$ and $\mu = 4.0$.

shadowing distance ε is significantly smaller in case of Milstein scheme compared to EM scheme. Because we do not consider the reparameterization of time, the value of shadowing distance ε are not small with respect to the local error δ .

	EM		T=10.5	2*	
μ	c	α	δ	β	ε
0	325.1272	1.4603	1.0500e-3	8.7832e-4	0.6858
1.0	327.3332	13.2132	0.0324	0.0008	22.0674
2.0	358.8756	91.1213	0.0324	0.0009	29.1598
4.0	377.5459	178.7624	0.0324	0.0008	36.0488

Finally, the approximation results of shadowing obtained from the shadowing method is shown in Fig.2. The blue cycle indicate the shadowing orbits and the black stars represent the (ω, δ) -pseudo orbits. We choose 50 points in the computational result. It presents that there are the shadowing orbits in the proper neighbourhood of the (ω, δ) -pseudo orbits. Therefore, the theoretical results can give information on chaotic orbits by the existence of shadowing orbits and the computation of shadowing distances.

	Milstein		T=10.5	2*	
μ	c	α	δ	β	ε
0	325.1272	1.4603	1.0500e-3	8.7832e-4	0.6858
1.0	332.3267	10.1223	1.0500e-3	0.0009	0.7191
2.0	368.8352	89.5351	1.0500e-3	0.0010	0.9626
4.0	398.7845	139.0756	1.0500e-3	0.0009	1.1295

Table 1: Summaries of the parameters for SCS.




Fig. 2. Zoom in parts of the approximation shadowing trajectory on SCS.

5.3 Compare with results of other SDEs

5.3.1 Results on stochastic Lorenz equations

We consider the Stratonovich SLEs

$$dX_t = f_0(X_t)dt + f_1(X_t) \circ dW_t(\omega), \quad X(0) = x_0 \in \mathbb{R}^3 \quad (19)$$

where the parameters are similar with SCS (16) and

$$f_0(X_t) = \begin{pmatrix} -\sigma x(t) + \sigma y(t) \\ \rho x(t) - y(t) - x(t)z(t) \\ -\beta z(t) + x(t)y(t) \end{pmatrix}, \quad f_1(X_t) = \begin{pmatrix} \mu x(t) \\ \mu y(t) \\ \mu z(t) \end{pmatrix}.$$

Here we choose the parameters $\sigma = 10, \rho = 28, \beta = 8/3$ and the initial value $x_0 = 0, y_0 = 1, z_0 = 0$. Table 2 contains some numerical results we obtained for SLEs.

Figures 3 and 4 show the pseudo orbits and the shadowing orbits for SLEs. As we can see, there are similar results as the results of SCS.

5.3.2 Results on stochastic lü equations

	EM		T=21.0	2*	
μ	c	α	δ	β	ε
0	468.9008	1.7147	1.0500e-3	0.0009	0.9883
0.5	451.2325	13.6754	0.0324	0.0010	30.1261
1.0	383.2516	84.3276	0.0324	0.0009	30.2991
1.3	356.3169	123.5653	0.0324	0.0008	31.0964

	Milstein		T=21.0	2*	
μ	c	α	δ	β	ε
0	468.9008	1.7147	1.0500e-3	0.0009	0.9883
0.5	458.3582	12.4722	1.0500e-3	0.0010	0.9887
1.0	417.7546	67.9523	1.0500e-3	0.0010	1.0199
1.3	378.6774	129.7812	1.0500e-3	0.0009	1.0678

Table 2: Summaries of the parameters for SLEs.

We consider the Stratonovich SLEs [7]

$$dX_t = f_0(X_t)dt + f_1(X_t) \circ dW_t(\omega), \quad X(0) = x_0 \in \mathbb{R}^3 \quad (20)$$

where the parameters are similar with SCS (16) and

$$f_0(X_t) = \begin{pmatrix} -\sigma x(t) + \sigma y(t) \\ \rho y(t) - x(t)z(t) \\ -\beta z(t) + x(t)y(t) \end{pmatrix}, \quad f_1(X_t) = \begin{pmatrix} \mu x(t) \\ \mu y(t) \\ \mu z(t) \end{pmatrix}.$$

Here we choose the parameters $\sigma = 36$, $\rho = 15$, $\beta = 3$ and the initial value $x_0 = 0$, $y_0 = 1$, $z_0 = 0$. Table 3 contains the numerical results we obtained for SLEs.

	EM		T=10.5	2*	
μ	c	α	δ	β	ε
0	456.1289	1.4728	1.0500e-3	0.0008	0.9609
0.5	449.8121	10.6576	0.0324	0.0009	29.8384
1.0	399.5433	75.8756	0.0324	0.0010	30.8071
1.3	366.7623	110.7654	0.0324	0.0010	30.9438

Figures 5 and 6 show the pseudo orbits and the shadowing orbits for SLEs. As we can see, there are similar results as the result of SCS.




Fig.3. Pseudo orbits for different sizes of noise for $\mu = 0, \mu = 0.5, \mu = 1.0$, and $\mu = 1.3$.




Fig. 4. Zoom in parts of the approximation shadowing trajectory on the SLEs.

As can be seen from these numerical results, there is an explicit dependent relationship between the shadowing distance and the local error, and there exists a true orbit in the appropriate neighbourhood of a (ω, δ) -pseudo orbit of SDE (16), (19) and (20), respectively. The numerically detected behavior of the system indeed reflects its real dynamical behavior which can be described more accurately.

5.4 The features of my device

And the features of my device are as follows, in which the computations are performed.

6. CONCLUSION

The main result of this paper is the numerical implementation of shadowing theorem for finite time

	Milstein		T=10.5	2*	
μ	c	α	δ	β	ε
0	456.1289	1.4728	1.0500e-3	0.0008	0.9609
0.5	448.4532	10.3534	1.0500e-3	0.0010	0.9635
1.0	388.3222	74.7751	1.0500e-3	0.0009	0.9725
1.3	365.8712	109.9121	1.0500e-3	0.0010	0.9991

Table 3: Summaries of the parameters for SLuEs.




Fig. 5. Pseudo orbits for different sizes of noise for $\mu = 0, \mu = 0.5, \mu = 1.0$, and $\mu = 1.3$.




Fig.6. Zoom in parts of the approximation shadowing trajectory on the SLuEs.

My device	Precision	RAM	Processor speed
Lenovo-PC	64-bit operating systems	6.00GB	2.20GHz

Table 4: The features of my device.

of SDEs. It focuses on the mathematical approaches to estimate the shadowing distance in the neighborhood of a (ω, δ) -pseudo orbit. The results show that the methods are effective and the numerical experiments are performed and match the results of theoretical analysis. More simple and high-performance methods will be shown in our further work.

ACKNOWLEDGEMENT

All data in this manuscript is available. Authors would like to express their gratitude to Prof. Jinqiao Duan, Prof. Xiaofan Li, Prof. Xiangdong Xie and Prof. Jialin Hong for their helpful discussions. The first author also thanks Prof. J. Duan and Prof. X. Li for their friendly reception during his visit (2019-2020) in Illinois Institute of Technology. This work is supported by NSFC (No. 61841302 and 11771449). This work is also supported by the Science Research Projection in the Education Department of Fujian Province, No. JT180122, Education Reform Fund of Fujian Agriculture and Forestry University, No. 111418136, the Natural Science Foundation of Fujian Province, No. 2015J01019, the Fundamental Research Funds for the Central Universities, HUST, No. 2016YXMS226, and China Scholarship Council, CSC No. 201907870004.

REFERENCES

1. L. Arnold, *Random dynamical systems*, Springer, 2003.
2. L. Arnold and B. Schmalfuss, Lyapunov's second method for random dynamical systems, *J. Differential Equations*, **177**(1) (2001), 235-265.
3. Y. Chen, H. Gao and C. Sun, The stochastic fractional power dissipative equations in any dimension and applications, *J. Math. Anal. Appl.*, **425** (2015), 1240-1256.
4. J. Duan, *An introduction to stochastic dynamics*, Cambridge University Press, 2015.
5. A. Fakhari and A. Golmakani, Shadowing properties of random hyperbolic sets, *Int. J. Math.*, **23** (2012), 1250025.
6. G. H. Golub and C. F. Van Loan, *Matrix computations*, 4th edition, The Johns Hopkins University Press, 2013.
7. J. Lü and G. Chen, A new chaotic attractor coined, *International journal bifurcation and chaos*, **12**(3) (2002), 659-661.
8. L. Li, P. Lin, X. Si, and L. Zheng, A numerical study for multiple solutions of a singular boundary value problem arising from laminar flow in a porous pipe with moving wall, *J. Comput. Appl. Math.*, **313** (2017), 536-549.

9. G. Milstein, *Numerical integration of stochastic differential equations*, Kluwer Academic Publishers, 1995.
10. T. Tél and M. Gruiz, *Chaotic dynamics: A introduction based on classical mechanics*, Cambridge University Press, New York, 2006.
11. K. J. Palmer, *Shadowing in dynamical systems, theory and applications*, Kluwer Academic Publishers, 2000.
12. S. Y. Pilyugin, *Shadowing in dynamical systems*, Springer, 1999.
13. D. Todorov, *Stochastic shadowing and stochastic stability*, arXiv:1411.7604v1.
14. P. Wang, A-stable Runge-Kutta methods for stiff stochastic differential equations with multiplicative noise, *Comput. and Appl. Math.*, (2014) 1-20.
15. T. Wang, Maximum error bound of a linearized difference scheme for coupled nonlinear Schrodinger equation, *J. Comp. Appl. Math.*, **235** (2011), 4237-4250.
16. X. Wang, J. Duan, X. Li and Y. Luan, Numerical methods for the mean exit time and escape probability of two-dimensional stochastic dynamical systems with non-Gaussian noises, *Appl. Math. Comput.*, **258** (2015), 282-295.
17. X. Xie and Q. Zhan, Uniqueness of limit cycles for a class of cubic system with an invariant straight line, *Nonlinear Anal. TMA*, **70** (2009), 4217-4225.
18. Q. Zhan, Mean-square numerical approximations to random periodic solutions of stochastic differential equations, *Adv. in Diff. Eqs.*, **292** (2015), 1-17.
19. Q. Zhan, Shadowing orbits of stochastic differential equations, *J. Nonlinear Sci. Appl.*, **9** (2016), 2006-2018.
20. Q. Zhan and Y. Li, Stochastic shadowing analysis of a class of stochastic differential equations, *J. Nonlinear Sci. Appl.*, **10** (2017), 5552-5565.
21. Q. Zhan and X. Xie, Numerical study of stochastic periodic Lipschitz shadowing of stochastic differential equations, *Discrete Dynamics in Nature and Society*, (2018), Article ID 1967508, 10 pages.
22. Q. Zhan, Shadowing orbits of a class of random differential equations, *Applied Numerical Mathematics*, **136**(1) (2019), 206-214.