

Skordalakes E and Berger J M (2003) Structure of the Rho transcription terminator: mechanism of mRNA recognition and helicase loading *Cell* **114**:135–46.

Steiner T, Kaiser J T, Marinković S, Huber R and Wahl MC (2002) Crystal structures of transcription factor NusG in light of its nucleic acid and protein-binding activities *EMBO J* **21**:4641–53.

Steinmetz E J and Platt T (1994) Evidence supporting a tethered tracking model for helicase activity of *Escherichia coli* Rho factor *Proc. Natl. Acad. Sci. U.S.A* **91**:1401–5.

Sullivan S L and Gottesman M E (1992) Requirement for *E. coli* NusG protein in factor-dependent transcription termination *Cell* **68**:989–94.

Thomsen N D and Berger J M (2009) Running in reverse: the structural basis for translocation polarity in hexameric helicases *Cell* **139**:523–34.

Toulokhonov I, Artsimovitch I and Landick R (2001) Allosteric control of RNA polymerase by a site that contacts nascent RNA hairpins *Science* **292**:730–33.

Valabhoju V, Agrawal S and Sen R (2016) Molecular basis of NusG-mediated regulation of Rho-dependent transcription termination in bacteria *J Biol Chem* **291**:22386–403.

von Hippel P H and Yager T D (1992) The elongation-termination decision in transcription *Science* **255**:809–12.

Washburn R S, Gottesman M E. (2011) Transcription termination maintains chromosome integrity *Proc Natl Acad Sci U.S.A* **108**:791–97.

Wilson K S and von Hippel P H (1994) Stability of *Escherichia coli* transcription complexes near an intrinsic terminator *J Mol Biol* **244**:36–51.

Accepted Version