

## **Wadia Institute of Himalayan Geology**

KISHOR KUMAR and PRADEEP SRIVASTAVA

*Wadia Institute of Himalayan Geology, 33, General Mahado Singh Road, Dehradun 248 001, India*

Wadia Institute of Himalayan Geology (WIHG) is an autonomous Institute of the Department of Science and Technology, Government of India. The mandate of this institution is to

- (a) Carry out research towards the development of new concepts and models for the geodynamic evolution of the Himalaya.
- (b) Develop exploration strategies and management of the natural reserves like minerals, industrial material, glaciers and water.
- (c) Plan and coordinate research towards understanding physical processes causing natural disasters, evaluate their hazard potential and suggest remedial measures.
- (d) Serve as national reference center for the database on the Himalayan Geology.

Research, in this institute, is broadly grouped into following thrust areas:

1. Himtransects
2. Climate-Tectonics Interaction
3. Biodiversity-Environment Linkage
4. Sustainable Natural Resources
5. Real Time Geology to Society: Coping with Natural Hazards.

These studies are performed at various platforms. For example, preparation of geological maps, sedimentological and structural mapping of the sequences, collection of data on microseismicity, and GPS studies are performed in field. Microscopic and geochemical chronological studies and data interpretation are done in laboratories. The field and laboratory data are integrated in order to understand various aspects of the evolution of the Himalaya and its active tectonics. Some important thrust areas where the institute has focused during the last five years are highlighted below:

### **Himtransects**

The semantic approach towards the Himalaya is the development of geodynamic model of the orogenic belt that can not only explain the archetype of the collision tectonics but can also trace the metamorphic history, constrain processes controlling high rate of erosion and exhumation, extent and nature of crustal shortening, kinematics of the crustal deformation, effects of superposed deformation as well as can provide clues to the space-time distribution of seismicity. The co-ordinated studies are in process along the HIMPROBE (a lithospheric geotransect across the Indian NW Himalaya) project in NW Himalaya. This transect has the advantage over the INDEPTH experiment as one single transect cuts across the entire Himalayan orogen from the Main Frontal Thrust to the Indus Suture Zone and all the essential facilities/expertise are available under one roof. The extended profile encompassing the Siwalik Range will help to trace the effects of post collision superposed deformation in shaping the oblique ramps. A similar integrated geotransect has also been initiated in the Eastern Himalayan Syntaxis region that remained largely unexplored and where the proposed mechanism of perpendicular continent-collision applicable along the Himalayan arc may not hold well because of the observed oblique subduction in the NE Himalaya. The challenge lies in the fact that geotransect in this region would constrain the role of collision, extension as well as shear tectonics. Below mentioned are major results accomplished under this thrust area:

- \* First ever passive seismology experiment carried out around the Eastern Himalayan Syntaxis shows presence of Moho at a depth of 47 km beneath the Brahmaputra Valley that gradually increases towards east and attains a depth of 62 km at the Suture zone.
- \* Petrochemical studies of granitoid rocks from the eastern part of Kumaun region suggest that the leading edge of India represented an active arc during the Late Paleoproterozoic time.

- \* The intra-crustal low velocity zone (IC-LVZ) beneath Ladakh within a depth range of 15-35 km has been identified. The presence of IC-LVZ and high Poisson's ratio in the crust signifies the presence of partial melt or aqueous fluids or both.
- \* The crustal growth in the Shyok-Darbuk corridor of NE Ladakh can be explained by two stage melting process involving the generation of orthogneiss with mafic enclaves (pre-collision), and temporal evolution towards increasingly more evolved felsic magmas during reworking processes (post-collision).
- \* Geochemical volume strain and elemental mobility studies on the mylonitized and extremely deformed phyllonitic quartzite along the MCT in the Bhagirathi Valley predict a very high fluid/rock ratio during development of the MCT zone.
- \* Minamiite, a Ca-bearing hydrous sulphate mineral, is reported for the first time in India from the Deccan Volcanic Province at Matanumadh (Kachchh, Gujarat). Hydrous sulphates are abundant on the planet Mars but not common on Earth because they form under extreme conditions. Therefore, these can serve as a potential Earth analogue for the Mars and the Martian conditions.
- \* Orientation patterns of magnetic susceptibility axes from AMS study of soft sediments from a trenched fault zone across the Himalayan Frontal Thrust (HFT), western Himalaya helped to identifying co-seismic structures in the Late Holocene surface sediments.
- \* Analysis of the geometry and folding pattern within the Chamba Thrust Sheet reveals that regional folds were developed during the southward translation of the Chamba Thrust Sheet. The structural mapping of the area between the Chamba and Tandi synclines further reveals that these synclines are ultimately connected with each other by the two anticlines and with a very open undulating syncline in between. The described structures fit well into asymmetric box fold geometry.
- \* The preliminary results of Geothermal springs and associated river waters collected from the Main Central Thrust (MCT) zone of the Himalaya show that there is a strong variation from -8.5‰ to +4.0‰ (PDB) indicating the varied sources of their origin.
- \* The temporal correlation of major suture zones with oceanic geoid anomaly revealed that the increase of geoid anomaly after 90 Ma in the vicinity of the Ninety East Ridge represents a phase of crustal

accumulation followed by the formation of Shyok suture.

### Climate-Tectonics Interaction

Recent researches have shown that tectonic deformation, climate and life interact in ways that are sometimes profound yet not very obvious. This has implications in terms of 1) the development of compressional orogens that is strongly influenced by climate where the extent of erosion determines the short and long term crustal deformation, 2) global geochemical processes and carbon and oxygen cycles that are soundly subjected by tectonics and climate change, and 3) Floral and faunal evolution that has strong bearing with climato-tectonic conditions. Therefore, integration of studies related to a) climate change, b) crustal deformation and its surface manifestations, c) surface processes in terms of geochemical changes, and d) floral and faunal responses to changing global conditions are likely to yield provocative results in climatic, biological, and tectonic interaction research in future.

In the Himalaya, it will be a challenging program to investigate whether the continental scale changes result from a) change in the mode and intensity of tectonic deformation, b) synchronous regional or global change of climate conditions, c) the dynamic evolution of the Himalayan drainage system, or a combination of all these processes. Therefore, most felt need of the contemporary research is to understand that how climate and orogenic organization of Indian subcontinent has responded to 1) continental tectonics at long term time scales of  $10^7$ - $10^5$  yrs since 60 Ma, 2) Regional crustal deformations at short time scales of  $10^2$ - $10^3$  yrs. Below are some recent findings under this thrust area:

- \* Two generations of alluvial fans, younger (23 to 10 ka) in the northwest and older (36 to 29 ka) in the southeast have been identified in the Soan Dun, which record the coupling of climate and tectonics. The uplifted Late Quaternary deposit, similar in age as the top of the older fan sequence, in the hanging wall of the Soan Thrust indicates tectonic activity north of the Himalayan Frontal Thrust (HFT), and suggests out-of-sequence thrusting during the Late Pleistocene.
- \* The Mio-Pliocene Middle Siwalik Subgroup of the Ranital-Kangra section of the Kangra sub-basin is sandstone dominated succession. It has been observed that at around 100 to 125 ka, it is interceded with a floodplain sequence of 1 to 3 m thick mudstone, which at places has also an abnormal thickness of >30 m. At basin scale, this cyclic pattern is controlled by extrinsic process; however, development of abnormally thick floodplain deposits

is controlled by intrinsic process due to basin tilting along the basin bound transverse fault.

- \* Sedimentation pattern of the Siwalik succession between 12 to 0.5 Ma, constrained through the reversal magneto-stratigraphy, exhibits two major phases of uplift at ~10 and ~5.5 Ma. The fluvial architecture and isotopic signatures, marking the intensification of the monsoon and accompanying changes in the drainage pattern show unambiguous evidence of climatic change triggered by the tectonic uplift, which was also responsible for re-organization of drainage pattern at ~10 Ma, ~5 Ma and 1.77 Ma in the foreland basin.
- \* The multi-parameter paleoclimate records of lacustrine sediments underlying the Chandra Peat deposit in Lahul and Spiti area have been precisely documented for the first time, as the Younger Dryas cold event in the NW Himalaya. Pollen data and mineral magnetic record indicating significant wet and warm conditions before 12880 cal yr BP, point towards the Ållerød interstadial climate. This cold/dry climate further intensified until 11640 cal yr BP and terminated with gradual reappearance of local as well as regional flora.
- \* Mineral magnetic studies from a proglacial lake of Chorabari suggest that variation in magnetic parameters is probably controlled by transportation energy related to the activity of the glacier.
- \* Patsio paleolake deposit, near the Bara Lacha Pass has revealed that it was developed due to climate-induced landslide and resultant blockade of the Bhaga River ca. 8-7 ka BP. The termination of fluvial sedimentary sequence and sudden onset of the glacial lake deposit marks an abrupt climate shift towards significant warmer conditions around 6.3 ka BP, which correspond to significant Holocene global climatic event.
- \* The Ganga River system in the NW and Brahmaputra River in the NE Himalaya during the past 40 ka aggraded in two major phases in response of the two phased deglaciation in the Higher Himalaya. The incision of the river valley took place in response to increased monsoon ~11 ka; however, in the foreland the incision took place 3-4 ka later suggesting a time lag in the surficial responses to particular climatic pulse.

### **Biodiversity-Environment Linkage**

Evolution is an innovative science which explains the ways in which the life has advanced, and whose signatures are found well preserved in the fossil records. Numerous lines

of evidence reveal that life has changed through time. The biodiversity includes the variability within and between the species and ecosystems. In geological context, it includes diversity within and among fossil taxa, and among paleo-ecosystems. Major geological boundaries are delineated based on distinct biological extinctions and recoveries attributed to ecosystem changes. Detailed understanding about the biodiversity across the major geological boundaries (i.e. PC/C, P/T, K/T, P/E, etc.) in the Himalaya is important for stratigraphic correlation.

Himalaya provides an ideal natural laboratory for testing the evolutionary innovation, diversification, extinction and environmental linkage with various bioevents. The Himalayan sedimentary records grant great opportunity to understand the linkages between biodiversity and environment at different time scales. An integrated multidisciplinary approach involving detailed taxonomy, morphometry, biostratigraphy-sedimentology, stable isotope and sedimentary geochemistry and magneto-stratigraphy is aimed to arrive at meaningful goal on the biodiversity-environmental linkage. Below are the major achievements under this thrust area:

- \* The oxygen isotope composition ( $\delta^{18}\text{O}$ ) of fossilized skeletal elements of vertebrates such as teeth and bones of sharks and other fish and some terrestrial mammals from the sub-Himalayan and western Peninsular Indian successions show strong correspondence with respective animal habitats. The freshwater  $\delta^{18}\text{O}$  values seem to suggest that monsoon like precipitation may have been active even during the Eocene-Oligocene period.
- \* Fossils bones and dentitions of rodents, primates, artiodactyls and other land-dwelling mammals recovered from the horizons corresponding to Early Eocene Climatic Optimum (EECO) show a rather close similarity with the contemporaneous fauna from Europe indicating faunal connection with that continent during the Early Eocene (~53 million year ago). This has opened a new debate regarding dispersal of smaller mammals across Europe and the Indian subcontinent.
- \* Molecular study of fossil resin from the Late Cretaceous Mahadeo Formation of Meghalaya indicates that, the major pyrolysis products and methyl-esterified thermochemolysis products found in the resin are abietane and labdane type diterpenoids, with minor amount of sesquiterpenoids. The exclusive presence of both labdane and abietane diterpenoids and the lack of phenolic terpenoids seem to suggest that, the Cretaceous resin was derived from Pinaceae (pine family) conifers.

- \* Nannofossils have been recovered from the Upper Bhuban Formation, Mizoram, northeast India, and probably it is for the first time that nannofossils from the Surma basin are recorded. The nannofossil assemblage suggests a hemipelagic depositional setting for the Upper Bhuban Formation of Mizoram.
- \* The identification and study of ancestral form of the whales (Cetacea) from a huge faunal collection from the Kalakot area, Jammu and Kashmir demonstrates that whales evolved from an aquatic group of mammals living in freshwater bodies rather than on land, necessitating the revisions in the views held so far that whales had land-dwelling ancestors.
- \* The Muth Formation in Pin valley has yielded an ichnofossil assemblage represented by a variety of ichnogenera, which show intertidal or littoral settings having a soft and sandy substrate and presence of annelid worms indicative of deep marine environment.
- \* The mineralogical, textural and geochemical characteristics of high-silica rhyolitic tuff breccia at the base of the late Paleocene-Middle Eocene Subathu Formation around Kalakot in Jammu and Kashmir provide a stratigraphic evidence of the earliest collision between the Indian and the Asian plates. The breccia, therefore, represent as an important marker horizon in the foreland stratigraphy.
- \* The black shales in the early Eocene part of the Subathu Formation in NW Himalaya occur in a relatively carbonate dominant sequence rich in organic matter and poor in carbonates as well as in fauna. They are lithologic expressions of hyperthermal event. An inverse relation between  $\text{CaCO}_3$  and TOC contents is observed in the black shales. This is likely in response to variations in seawater temperature, dissolved oxygen and/or pH.
- \* Organic remains from the Dharamsala Group of Kangra Valley indicate that the beginning of orogeny was apparently a key driver for resurgence of life on southern flank of the Himalaya during the early Miocene time. The early Eocene life known from the older sequence of the area was plausibly wiped out by global sea level fall as a precursor to the beginning of Ice Age Earth.
- \* Evidence of well preserved uvigerinid foraminifera and pteropods in the Tertiary horizons of NE India signifies that the hydrocarbon producing Upper Disang Formation was deposited at a depth of ~500 m, well above the aragonite compensation depth (ACD) in an open marine basin, information crucial in oil exploratory activity in the region.

- \* Neoproterozoic Organic Walled Microfossils (OWMs) including silicified cyanobacteria and acritarchs have been reported from the Infrakrol and lower Krol formations of outer carbonate belt and Gangolihat Formation of Inner Carbonate belt. Sponge spicules of similar age have also been documented from both outer as well as inner carbonate belts, bringing down their known range.

### Sustainable Natural Resources

Resources form the backbone of every economy. The very use of resources and their transformation help in building the capital stocks that add to the wealth of present and future generations. In this context, the Himalaya is a special region in the mountain ecosystems of the world. This geodynamically active region is not only important from the viewpoint of climate and as a provider of life, water resource to large part of the subcontinent, but it also houses a variety of flora, fauna, and human communities. Despite the abundance of natural resources, most of its people are living on the edge on subsistence level. A judicious management and study of glaciers, streams, biota and mineral resources in totality is needed for sustained development of economic, ecological and socio-cultural necessities. It is therefore essential to understand the various conditions and processes, which control and affect the occurrence of natural resources particularly, water (glaciers, streams, and groundwater) and mineral. It is therefore proposed to address the aforementioned themes on natural resources, which also have immense societal importance and are essential to accelerate the development activities in our country. Some of the results achieved from this research programme are mentioned below:

- \* Field evidences, on the basis of loops of lateral and terminal moraines suggest that there were four phases of advances and retreat of the Dokriani-Bamak glacier over the last ~315 years.
- \* The dates calculated by lichenometry of different loops of deglaciation indicate the influence of climate change on the glaciers and difference in value suggests the effect of orientations of the two glaciers. The north facing Dokriani glacier in the Garhwal Himalaya indicates Little Ice Age maximum during the AD 1692, whereas the south facing Chorabari glacier shows its maximum during the AD 1748.
- \* Treeline is one of the natural and important climate markers in the high altitude mountain ecosystems. Investigations carried out in the Chorabari and Dokriani glaciers suggest that the treeline has shifted towards higher altitudes at the rate of 10.21m/year in Chorabari and 1.70m/year in Dokriani glacier valleys with in a period of 47 years.

- \* 'Stannoidite', an unusual sulphide of Cu, Fe, Sn and Zn within the chalcopyrite is reported for the first time from the eastern Kumaun region, which suggest the genesis of polymetallic sulphide mineralization due to cooling of high temperature and sulphur fugacity hydrothermal solution.
- \* The observed geochemistry of Pinjaur Dun sediment samples probably reflects both the mechanical dispersion/erosion of catchment's host rocks as well as the hydromorphic (aqueous phase) transport and transfer of REE from the aqueous phase to the stream sediments. The high  $(La/Yb)_N$  ratios along with the major and trace element chemistry in the samples suggest dominance of a felsic provenance and deposition in an environment dominated by a particulate terrigenous source.
- \* Geo-accumulation index (I-geo), applied for the purpose of environmental evaluation of heavy metals distribution in Pinjaur Dun area, indicated that leaching of metals from garbage and industrial sewage (effects of anthropogenic activities) has led to enhancement of metal concentration (Pb, Zn and Cu).
- \* The representative carbon isotope analysis ( $\delta^{13}C$ ) of Almora graphite suggested that the graphite has crystallized from biogenic carbon during the metamorphism of the host sediments.

### Real Time Geology to Society: Coping with Natural Hazards

The Himalaya is an abode of natural hazards. In addition to the seismic hazard, the mountain belt recurrently witnesses landslides, cloudbursts, flash floods, snow avalanches, glacier lake outburst floods, etc. The frequent occurrences of natural hazards pose a continual threat to the safety of the people inhabiting in and adjacent to this gigantic mountain system. It is therefore essential to enhance our knowledge base on various physical and geodynamic processes causing natural hazards possibly in a real time mode for developing and implementing suitable mitigation measures. This line of fundamental research with serious societal implications forms the central focus of the mission mode project. As a preventive measure to minimize the damage scenario in the event of a large earthquake, microzonation maps depicting dominant frequencies in propagating seismic waves and their amplification by near surface sedimentary column are proposed for major townships and industrial belt, including possible sites of hydro-electric dams. The research programme focusing on

landslides, cloud bursts, landslide lake outburst floods and glacial lake outburst floods will establish the kinematic model to identify interactive internal causative factors, which would help in planning of mitigation strategies. The highlights of results under this thrust area are given below:

- \* A new program combining powerful InSAR and on-going GPS campaign surveys around the reservoir was taken up to constrain better the uplift/subsidence associated with annual loading/unloading of the reservoir. Long term monitoring of GPS, total magnetic field intensity and radon emission are also in progress at 14 stable bench marks established around the Tehri reservoir.
- \* The recent trench excavation survey carried out across an active fault system in the Quaternary deposits of the Pinjaur Dun demonstrates a 2 meter vertical displacement substantiating our previous reports of large magnitude earthquakes rupturing the frontal Himalaya.
- \* Continuous radon monitoring in a 68 m deep borehole in the Garhwal Himalaya, and the abnormalities observed in radon emanation and other parameters in the borehole have shown to be precursory signature for the July 23, 2007 Kharsali earthquake of M4.9, that had an epicenter distance of 60 km.
- \* Seismic hazard analysis of the NW Himalaya carried out using probabilistic technique show high hazard potential zones to be centered around the Kashmir region (0.70g), Kangra region (0.50g), Garhwal region (0.50g) and Darchula region (0.50g) with intervening zones representing the low hazard zones (~0.25g) for 10% probability of exceedance in 50 years.
- \* A new methodology has been developed based on temporal gravity and solid earth deformation studies to understand the dynamics of alluvial pore space fluid in the Doon Valley, when it was subjected to distant large or great magnitude earthquakes.
- \* More recently, a moderate (M~6.9) earthquake which occurred in Sikkim, (NE India) on September 18, 2011 is being monitored by the WIHG with the help of eight broadband seismographs installed in and around the epicentral region. The studies are aimed at understanding the size and orientation of the fault plane responsible for the Sikkim Earthquake from the aftershock observations.

**Significant Publications 2007-08**

- Asthana, A.K.L. and Sah, M.P., 2007. Landslides and cloudburst in the Mandakini basin of Garhwal Himalaya. *Him. Geol.*, v. 28, p. 59-67.
- Azmi, R.J. and Joshi, D., 2007. Comment on paper "A Neoproterozoic geomagnetic field reversal from the Kurnool Group, India: implications for stratigraphic correlation and formation of Gondwana" by M.R. Goutham et al. *Jour. Geol. Soc. India*, v. 67, p. 221-233. *Jour. Geol. Soc. India*, v. 70, p. 876-878.
- Bajpai, S., Das, D.P., Kapur, V.V. Tiwari, B.N. and Srivastava, S.S., 2007. Early Eocene rodents (Mammalia) from Vastan Lignite Mine, Gujarat, Western India. *Geol. Mag.*, v. 22, p. 91-95.
- Banerjee, P., Pollitz, F. and Burgmann, R., 2007. Coseismic slip distribution of the 26 December 2004 Sumatra-Andaman and 28 March 2005 Nias earthquakes from GPS static offsets. *Bull. Seism. Soc. America*, v. 97, No.1A, pp. S1-S, doi: 10.1785/0120050609.
- Bartarya, S.K., Mazari, R.K. and Virdi, N.S., 2007. Bhimgoda slide of August 24, 1994 in the Siwalik rocks near Haridwar : a success story of landslide control measures. In *Natural Hazards, Spl. Publ. Ind. Geol. Cong.*, p. 155-167.
- Brook, G.A., Marais, E., Srivastava, P. and Jordan, T., 2007. Timing of lake-level changes in Etosha Pan, Namibia, since the middle Holocene from OSL ages of relict shorelines in the Okondeka region. *Quat. Internat.*, v. 175, p. 29-40.
- Chabak, S.K. and Sharma, P.K., 2007. Numerical Simulation of coupled wave Equations. *Internat. Rev. Pure and App. Math.*, v. 3, p. 81-91.
- Chakrapani, G., Mahanta, C., Routh, J., Magnus, M. and Khanna, P.P., 2008. Coupling between seismic activity and hydrogeochemistry at the Shillong Plateau, Northeastern India. *Pure and App. Geophys.*, v. 165, p. 45-61.
- Pant, C.C. and Paul, A., 2007. Recent trends in seismicity of Uttarakhand. *Jour. Geol. Soc. India*, v. 70, p. 619-626.
- Choubey, V.M., Mukherjee, P.K., Bajwa, B.S. and Walia, V., 2007. Geological and tectonic influence on water-soil radon relationship in Mandi-Manali area, Himachal Himalaya. *Environ. Geol.*, v. 52, p. 1163-1171.
- Choudhury, S., Rajpal, H. and Saraf, A.K., 2007. Mapping and forecasting of North Indian winter fog: an application of spatial technologies. *Internat. Jour. Remote Sensing*, v. 28, p. 3649-3663.
- Dobhal, D.P., Gergan, J.T. and Thayyen, R.J., 2007. Mass balance and snout recession measurement (1991-2000) of Dokriani glacier Garhwal Himalaya India, *Internat. Hydro. Prog.. Technical Documents in Hydrology*, v. 80, p. 53-63.
- Dobhal, D.P., Gergan, J.T. and Thayyen, R.J., 2008. Mass balance studies of Dokriani glacier from 1992 to 2000, Garhwal Himalaya, India. *Bull. Glaciological Res.*, v. 25, p. 9-17.
- Dubey, A.K. and Bhakuni, S.S., 2008. Neotectonic stresses and active tectonics in the Garhwal Inner Lesser Himalaya: anisotropy of magnetic susceptibility studies. *Him. Geol.*, v. 29, p. 35-47.
- Goel, O.P. and Bhakuni, S.S., 2007. Petrochemistry and petrogenesis of the Great Himalayan gneisses and granites from Chamoli District, Kumaun-Garhwal Himalaya. In: *Himalaya (Geological Aspect P.S. Saklani., Ed.)*, Satish Publishing House, New Delhi, v. 51-101.
- Gupta, V. and Ahmed, I., 2007. The effect of pH of water and mineralogical properties on the slake durability (degradability) of different rocks from the Lesser Himalaya, India. *Eng. Geol.*, v. 95 p. 79-87.
- Gupta, V. and Ahmed, I., 2007. Geotechnical characteristics of Surabhi Resort landslide in Mussoorie, Garhwal Himalaya, India. *Him. Geol.*, v. 28, p. 21-32.
- Gupta, V., Bist, K.S., Sharma, B. and Asthana, A.K.L., 2007. Varunavat landslide in the Bhagirathi valley, Garhwal Himalaya: its causes and risk assessment. In: *Natural Hazards, Spl. Publication, Ind. Geol. Cong.*, p. 97-111.
- Gururajan, N.S. and Choudhuri, B.K., 2007. Geochemistry of the Trans Himalayan Lohit Plutonic Complex, Eastern Arunachal Pradesh. *Jour. Geol. Soc. India*, v. 70, p. 17-33.
- Jowhar, T.N., 2007. Significance of chemical zoning in garnets. *Indian Mineralogist*, v. 41, p. 200-210.
- Jowhar, T.N., 2007. Pressure-Temperature paths from zoned metamorphic garnets. *Ind. Jour. Geochem.*, v. 22, p. 419-429.
- Jayangondaperumal, R. and Thakur, V.C., 2008. Kinematics of secondary surface fractures on southeastern extension of rupture zone, and 8 October 2005, Muzafarabad earthquake. *Tectonophysics*, v. 446, p. 61-76.
- Kamal and Mundepe, A.K., 2007. Site response studies in Dehradun city: first step towards seismic microzonation. In: *Natural Hazards, Sp. Pub., Ind. Geol. Cong.*, p. 175-181.
- Kapur, V., Bajpai S., Das, D.P. and Tiwari B.N., 2007. Early Eocene Primate (Mammalia) from Vastan Lignite Mine, District Surat (Gujarat), Western India. *Jour. Palaeo. Soc. India*, v. 52, p. 231-234.
- Kumar, K., Rana, R.S. and Singh, H., 2007. Fishes of the Khuiala Formation (Early Eocene) of the Jaisalmer Basin, Western Rajasthan, India. *Curr. Sci.*, v. 93, p. 553-559.
- Kumar, S., Chander, R. and Khattri, K.N., 2007. Upper mantle velocity structure in the NW Himalaya: Hindu Kush to Garhwal region from travel time studies of deep Hindu Kush earthquakes. *Edu. Res. and Review*, v. 2, p. 302-314.
- Kumar, S., Weskowsky, S.G., Rockwell, T.K., Briggs, R.W., Thakur, V.C. and Perumal, R.J., 2007. Paleoseismic evidence of great surface rupture earthquakes along the Indian Himalaya. *Jour. Geophys. Res.*, v. iii, B03304, 10., 1029/2004JB00309.
- Lokho, K. and Raju, D.S.N., 2007. Langhian (early Middle Miocene) foraminiferal assemblage from Bhuban Formation, Mizoram, NE India. *Jour. Geol. Soc. India*, v.70 (6), p. 933-938.
- Lokho, K. and Kumar, K., 2008. Fossil pteropods (Thecosomata, holoplanktonic Mollusca) from the Eocene of Assam-Arakan Basin, north-eastern India. *Cur. Sci.*, v. 94 (5), p. 647-652.
- Luirei, K. and Bhakuni, S.S., 2007. Soft sediment deformational structures in the lacustrine deposits of Ziro valley, Lesser Himalaya, Arunachal Pradesh. *Jour. Geol. Soc. India*, v. 70, p. 73-78.
- Luirei, K. and Bhakuni, S.S., 2007. Landslides along frontal part of Eastern Himalaya in East Siang and Lower Dibang districts, Arunachal Pradesh, India. *Jour. Geol. Soc. India*, v. 71, p. 321-330.
- Mahajan, A.K. and Ghosh, G.K., 2007. Statistical analysis of earthquake data from Northwest Himalayan region and its implication for

- seismic hazards. In Natural Hazards (Varma, O.P., Mahajan, A.K., Gupta, V. Eds), Sp. Pub., Ind. Geol. Cong., p. 45-56.
- Mahajan, A.K., Rob J. Sporry, Champati, P.K., Ranjan, Rajiv, Slob, S. and Cees J. van Westen, 2007. Methodology for site response studies using Multi-channel Analysis of Surface Wave (MASW) technique in Dehradun city. *Curr. Sci.*, v. 92, p. 945-955.
- Mahajan, A.K., Siefko Slob, Rajiv Ranjan, Rob Sporry, Champati, P.K. and Cees J. van Westen, 2007. Seismic Microzonation of Dehradun city using Geophysical and Geotechnical Characteristics in the upper 30-meters of soil column. *Jour. Seismology*, v. 11, p. 335-370.
- Mazari, R.K., 2007. Outline geomorphology of the upper Bhagirathi basin, Garhwal Himalaya. *Him. Geol.*, v. 28, p. 45-57.
- Misra, D.K., 2007. Evidences of Neotectonic activity along active faults in Arunachal Himalaya, NE India. *Him. Geol.*, v. 28, p. 75-78.
- Misra, D.K., 2007. Geomorphic features along the active faults in the Lohit and Dibang Valleys, Eastern Arunachal Pradesh, India. *Zeitschrift fur Geomorphologie N.F.*, v. 51(3), p. 327-336.
- Mondal, S.K., Sastry, R., Pachauri, A.K. and Gautam, P.K., 2008. High resolution 2D electrical resistivity tomography to characterize active Naitwar Bazar landslide, Garhwal Himalaya, India. *Curr. Sci.*, v. 94, p. 871-875.
- Mukherjee, P.K. Purohit, K.K. Saini, N.K. and Khanna, P.P., 2007. Geochemical landscape of Garhwal Lesser Himalaya based on active stream sediment geochemistry. *Ind. Jour. Geochem.*, v. 22, p. 79-89.
- Mundepi, A.K., 2007. A comparative study of site amplification by using horizontal to vertical spectral ratio (Nakamura technique) of ground ambient noise (GAN) with earthquakes at various lithounits in NW Himalaya. *Him. Geol.*, v. 28, p. 69-74.
- Mundepi, A.K., Kamal and Paul, A., 2007. Estimation of site amplification at various lithounits in NW Himalaya using horizontal to vertical ratio. *Jour. Geol. Soc. India*, v. 70, p. 605-609.
- Mundepi, A.K., 2008. Site response studies using horizontal to vertical ratio (H/V) in the city of Chandigarh, North India. *Him. Geol.*, v. 29, p. 87-93.
- Nanda, A.C. and Sehgal, R.K., 2007. Geology and mammalian faunas of the Siwalik Group of Northwestern Himalaya: recent investigations. In *Human Origins, Genome & People of India Genomic, Palaeontological and Archaeological Perspectives* (Sankhyan, A.R., Rao, V.R. Eds), Allied Publishers Pvt. Ltd, New Delhi, p. 77-90.
- Narasimhon, C.L. Patil, S.K. and Arora, B.R., 2008. A Palaeomagnetic perspective on the flows and intrusives of Kachchh Rift Basin and its implication on Reunion Plume-Indian Plate interaction. *In: Indian Dykes: Geochemistry, Geophysics and Geochronology: (Srivastava, R., Shivaji, C. & Chalapathi Rao, N.V. Eds)*, Narosa Publications, New Delhi, p. 159-180.
- Negi, P.S., 2007. Alien flora of Doon Valley, North-West Himalaya. *Curr. Sci.*, v. 92(7), p. 968-978.
- Nummela, S., Thewissen, J.G.M., Bajpai, S., Hussain, S.T. and Kumar, K., 2007. Sound transmission in archaic and modern whales: anatomical adaptations for underwater hearing. *The Anatomical Record*, v. 290, p. 716-733.
- Panda, S.K., Choudhury, S., Saraf, A.K. and Das, J.D., 2007. MODIS land surface temperature data detects thermal anomaly preceding 08 October 2005 Kashmir earthquake. *Internat. Jour. Remote Sensing*, v. 28(20), p. 4587-4596.
- Pant, P.D., Kothiyari, G.C. and Luirei, K., 2007. Geomorphologic and geological investigation of neotectonic activity of Saryu River Fault (SRF), a part of North Almora Thrust (NAT) in Seraghat-Basoli area in Central Kumaun, Uttaranchal. *Jour. Geol. Soc. India*, v. 70, p. 815-823.
- Parcha, S.K., Sabnis, S.V. and Saraswati, P.K., 2007. Taxonomic application of classification and regression tree (CART) and random forests (RF): a case study of Middle Cambrian Trilobites. *Jour. Geol. Soc. India*, v. 70, p. 1033-1038.
- Patil, S.K. and Arora, B.R., 2008. Palaeomagnetic and rock magnetic studies on the intrusives from Raniganj Basin, Damodar Valley: linkage to Rajmahal Volcanics. In *Indian Dykes: Geochemistry, Geophysics and Geochronology: (Srivastava, R., Shivaji, C. & Chalapathi Rao, N.V. Eds)*, Narosa Publications, New Delhi, p. 511-526.
- Paul, A., Pant, C.C., Darmwal, G.S. and Pathak, V., 2007. Seismicity of Uttarakhand as recorded by Kumaun Digital Telemetered Seismic Network. *In: Natural Hazards Spl. Publication, Ind. Geol. Cong.*, p. 57-63.
- Paulsen, Timothy, S., Christie M. Demosthenous., Myrow, P.M., Thompson, K.R., Hughes, N.C. and Parcha, S. K., 2007. Palaeostratigraphic analysis of calcite twins across the Cambrian-Ordovician unconformity in the Tethyan Himalaya, Spiti and Zaskar valley regions. *India. Jour. Asian Earth Sci.*, v. 31(1), p. 44-54.
- Philip, G., 2007. Remote sensing data analysis for mapping active faults in northwestern part of Kangra valley, NW Himalaya, India. *Internat. Jour. Remote Sensing*, v. 28(21), p. 4745-4761.
- Pollitz, F.F., Banerjee, P., Kelly, Grijalva, Nagarajan and Burgmann, Roland, 2008. Effect of 3D viscoelastic structure on postseismic relaxation from the 2004 M=9.2. Sumatra earthquake. *Geophys. Jour. Internat.*, doi:10.1111/j.1365-246X.2007.03666.x
- Purohit, K.K., Mukherjee, P.K., Saini, N.K., Rathi, M.S. and Khanna, P.P., 2007. Geochemical dispersion pattern of some heavy metals in upper Alaknanda catchment: application in mineral exploration and its environmental implications. *Jour. Geol. Soc. India*, v. 70, p. 97-108.
- Rana, R.S., Kumar, K., Escarguel, G., Sahni, A., Rose, K.D., Smith, T., Singh, H. and Singh, L., 2008. An ailuravine rodent from the early Eocene Cambay formation at Vastan, western India, and its palaeobiogeographic implications. *Acta Palaeontologica Polonica*, v. 53 (1), p. 1-14.
- Rao, D.R. and Rai, H., 2007. Permian komatiites and associated basalts from the marine sediments of Chhongtash Formation, southeast Karakoram, Ladakh, India. *Mineral. Petrol.*, v. 91 (3-4), p. 171-189.
- Rao, D.R. and Rai, H., 2007. Are garnet-amphibolites and eclogites from Tso-Morari region, Ladakh evolved in a rift environment. *Deep Continental Studies-DST-Newsletter*, v. 17(2), p. 2-6.
- Rathi, G., Sangode, S.J. Kumar, R. and Ghosh, S.K., 2007. Magnetic fabrics under high-energy fluvial regime of the Himalayan Foreland Basin, NW Himalaya. *Curr. Sci.*, v. 92, p. 933-944.

- Sachan, H.K, Mukherjee, B.K. and Bodnar, R.J., 2007. Preservation of methane generated during serpentinization of upper mantle rocks: evidence from fluid inclusions in the Nidar ophiolite, Indus Suture Zone, Ladakh (India). *Earth Planet. Sci. Lett.*, v. 257, p. 47-59.
- Sachan, H.K, Mukherjee, B.K. and Ahmad, T., 2007. Brine-rich Hydrothermal fluid circulation in the upper level of Nidar Ophiolite sequence, Ladakh: evidences from fluid inclusions. *Jour. Geol. Soc. India*, v. 70, p. 780-786.
- Sangode, S.J. and Mazari, R.K., 2007. Late Quaternary environmental changes in the Trans Himalayan Spiti valley, India: results of mineral magnetic studies on Kioto palaeolake. *Him. Geol.*, v. 28(2), p. 1-9.
- Saraf, A.K., Sinha, S.T., Ghosh, P. and Choudhury, S., 2007. A new technique to remove false topographic perception phenomenon and its impacts in image interpretation. *Int. Jour. Remote Sensing*, v. 28, p. 811-821.
- Saraf, A.K., Choudhury, S., Panda, S.K., Dasgupta, S. and Rawat, V., 2007. Does a major earthquake precede a thermal anomaly? *Internat. Jour. Geoinformatics*, v. 3(3), p. 1-8.
- Sinha, R., Kumar, R., Sinha, S., Tandon S.K. and Gibling, M.R., 2007. Late Cenozoic fluvial deposits of India: an overview. *Quatern. Sci. Rev.*, v. 26, p. 2801-2822.
- Sinha, S., Islam, R., Ghosh, S. K., Kumar, R. and Sangode, S.J., 2007. Geochemistry of Neogene Siwalik mudstones along Punjab re-entrant, India: implications for source-area weathering, provenance and tectonic setting. *Curr. Sci.*, v. 92, p. 1103-1113.
- Singh, A.K., 2007. Chemical characteristics of alkaline basalt from the Abor volcanics of Arunachal Himalaya. *Jour. Geol. Soc. India*, v. 69, p. 189-1194.
- Singh, S.K. Rai, S.K. and Krishnaswami, S., 2008. Sr and Nd isotopes in river sediments from the Ganga Basin: Sediment provenance and spatial variability in physical erosion, *Jour. Geophys. Res.*, v. 113, F03006, doi: 10.1029/2007JF000909.
- Singh, T. and Virdi, N.S., 2007. Tectonic activity classes along the Nahan thrust (NT) in NW sub Himalaya. *Jour. Ind. Soc. Remote Sensing*, v. 35(3), p. 221-230.
- Siva Siddaiah, N., 2008. Berthierine-rich Ooidal Ironstone from the Late Paleocene- Middle Eocene Subathu Formation, Dogadda area, NW Himalaya and its Stratigraphic Significance. *Curr. Sci.*, v. 94(1), p. 123-127.
- Sundriyal, Y.P., Tripathi, J.K., Sati, S.P., Rawat, G.S. and Srivastava, P., 2007. Landslide dammed lakes in the Alaknanda basin (Lesser Himalaya): causes and implications. *Curr. Sci.*, 93, 568-574.
- Suresh, N., Bagati, T.N., Kumar, R. and Thakur, V.C., 2007. Evolution of Quaternary alluvial fans and terraces in the intramontane Pinjaur Dun, Sub-Himalaya, NW India: interaction between tectonics and climate change. *Sedimentology*, v. 54, p. 809-833.
- Tewari, V.C., 2007. The rise and decline of the Ediacaran biota: palaeobiological and stable isotopic evidence from the NW and NE Lesser Himalaya, India. *Geol. Soc. London, Special Publication*, v. 286, p. 77-101.
- Tewari, V.C., Stenni Barbara, Pugliese Nevio, Drobne Katica, Riccamboni Rodolfo and Dolence Tadej, 2007. Peritidal sedimentary depositional facies and carbon isotopic variation across K/T boundary carbonates from NW Adriatic platform. *Palaeogeogr. Palaeoclim. Palaeoecol.*, v. 255, p. 77-86.
- Thewissen, J.G.M. Cooper, L.N. Clementz, M.T. Bajpai, S. and Tiwari, B.N., 2007. Whales originated from aquatic artiodactyls in the Eocene epoch of India. *Nature*, v. 450(7173), p. 1190-1194.
- Tiwari, M., 2008, Additional Neoproterozoic sponge spicules from Gangolihat Dolomite, Kumaun Lesser Himalaya, India. *Him. Geol.*, v. 29, p. 49-55.
- Virdi, N.S. and Gupta, V., 2008. Hydrel projects in the Yamuna and Tons valleys, Dehradun. *Excursion Guide, Publ. CAS in Geology, Panjab University, Chandigarh*, p. 20.
- Yeats, R.S. and Thakur, V.C., 2008. Active faulting south of the Himalayan front: establishing a new plate boundary. *Tectonophysics*, v. 453, p. 63-73

## 2008-09

- Ahmad T., Tanaka T., Sachan, H.K, Asahara Y., Islam R. and Khanna P.P., 2008. Geochemical and isotopic constraints on the age and origin of the Nidar Ophiolitic Complex, Indus Suture Zone, Ladakh, India. *Tectonophysics*, v. 451, p. 206-224.
- Banerjee, P., Bürgmann, R., Nagarajan, B. and Apel, E., 2008. Intraplate deformation of the Indian subcontinent. *Geophys. Res. Lett.*, v. 35, L18301, doi:10.1029/2008GL035468.
- Devrani, U. and Dubey, A. K., 2008. Anisotropy of magnetic susceptibility and petrofabric studies in the Garhwal Synform, Outer Lesser Himalaya: evidence of pop-up klippen. *Island Arc*, v. 18, p. 428-443.
- Gupta V. and Sah, M.P., 2008. The Relationship between Main Central Thrust (MCT) and the Spatial Distribution of Mass Movement in the Satluj Valley, Northwestern Higher Himalaya, India, *Zeitschrift für Geomorphologie*, v. 52 (2), p. 169-179.
- Gupta V. and Sah, M.P., 2008. Impact of the Trans-Himalayan Landslide Lake Outburst Flood (LLOF) in the Satluj Catchment, Himachal Pradesh, India. *Natural Hazards*, v. 45, p. 379-390.
- Kumar Naresh, Sharma, J., Arora, B. R., and S., Mukhopadhyay, 2009. Seismotectonic model of the Kangra-Chamba sector of NW Himalaya: Constraints from joint hypocenter determination and focal mechanism. *Bull. Seismol. Soc. Amer.* v. 99(1), p. 95-109
- Lokho, K. and Kumar, K., 2008. Fossil pteropods (Thecosomata, holoplanktonic Mollusca) from the Eocene of Assam-Arakan Basin, north-eastern India. *Curr. Sci.*, v. 94 (5), p. 647-652.
- Mahajan, A.K., 2009. NEHRP soil classification and estimation of 1-D site effect of Dehradun fan deposits using shear wave velocity. *Eng. Geol.*, v. 104, p. 232-240.
- Mathur, N.S., Juyal, K.P. and Kumar, K., 2008. Biotic response to Cretaceous-Eocene tectonic events at the northern margin of the Indian Plate and the Indus Suture Zone, Ladakh Himalaya, India. *Jour. Palaeo. Soc. India*, v. 53 (1), p. 37-49.
- Mukherjee, P.K. and Gupta, P.K., 2008. Arbitrary scaling in ISOCON method of geochemical mass balance: an elevation of the graphical approach. *Geochem. Jour.*, v. 42, p. 247-253.
- Nanda, A.C., 2008. Comments on the Pinjor Mammalian Fauna of the Siwalik Group in relation to the post-Siwalik faunas of Peninsular India and Indo-Gangetic Plain. *Quarter. Internat.*, v. 192, p. 6-13.
- Patel, S.C. Ravi, S. Anilkumar, Y. Naik, A. Thakur, S.S. Pati, J.K. and Nayak, S.S. 2009. Mafic xenoliths in Proterozoic kimberlites from Eastern Dharwar Craton, India: mineralogy and P-T regime. *Jour. Asian Earth Sci.*, v. 34, p. 336-346.



- Ramola, R.C., Choubey, V.M., Negi, M.S., Yogesh Prasad and Ganesh Prasad., 2008. Radon occurrence in soil-gas and groundwater around an active landslide. *Radiation Measurements*, v. 43(1), p. 98-101.
- Ramola, R.C., Yogesh Prasad, Kumar, Sushil. and Choubey, V.M., 2008. Soil-gas radon as seismotectonic indicator in Garhwal Himalaya. *Jour. Appl. Rad. Isotopes*, v. 66(10), p. 1523-1530.
- Rana, R.S., Kumar, K., Escarguel, G., Sahni, A., Rose, K.D., Smith, T., Singh, H. and Singh, L., 2008. An ailuravine rodent from the early Eocene Cambay Formation at Vastan, western India, and Its palaeobiogeographic implications. *Acta Palaeont. Polonica*, v. 53 (1), p. 1-14.
- Sangode, S. J., Kumaravel, V., Bloemendal. J. and Kumar. R., 2008. Effect of Burial and Compaction on Soil Magnetic Properties: Results from Soil-Paleosol sequences in the Himalayan Foreland, India. *Palaeogeogr. Palaeoclim. Palaeoecol.*, v. 267, p. 235-244.
- Saraf, A.K., Rawat, V., Banerjee, P., Choudhury, S., Panda, S.K., Dasgupta, S., and Das, J.D., 2008. Satellite Detection of earthquake thermal precursors in Iran. *Natural Hazards*, v. 47 (1), p. 119-135.
- Schopf, J.W., Tewari, V.C. and Kudryavtsev, A. B., 2008. Discovery of a new chert – permineralised microbiota in the Proterozoic Buxa Formation of the Ranjit Window, Sikkim, Northeast India, and its Astrobiological implicatios, *Astrobiology*, v. 8, p. 735- 746.
- Sharma, R. and Rameshwar Rao, D., 2008. Stannoidite and cubanite from Askot polymetallic sulphide ores, Kumaun Lesser Himalaya, India, *Current Science*, v. 95 (5), p. 527-531.
- Siddaiah, N.S. and Kumar, K., 2008. Tonstein (volcanic ash) from Late Paleocene-Middle Eocene sediments of the northwest Himalaya and its significance for the timing of initiation of India-Asia Collision. *Mem. Geol. Soc. India*, v. 72, p. 145-164.
- Singh, R.K.B., Gururajan, N.S. and Singh, A.K., 2008. Petrology and Geochemistry of Amphibolites of the Bomdila Group, Western Arunachal Himalaya. *Jour. App. Geochem.*, v. 10(2), p. 92-107.
- Srivastava, P., Bhakuni, S.S., Luirei, K., and Misra, D.K., 2009. Fluvial records from the Brahmaputra River exit, NE Himalaya: climate-tectonic interplay during Late Pleistocene-Holocene. *Jour. Quart. Sci.*, v. 24, p. 175-188.
- Srivastava, P. and Misra, D.K., 2008. Morpho-sedimentary records of active tectonics at the Kameng river exit, NE Himalaya, *Geomorphology*, v. 96, p. 187-198.
- Srivastava, P., Tripathi, J.K., Islam, R. and Jaiswal, M.K., 2008. Fashion and Phases of Late Pleistocene aggradation and incision in Alaknanda River, western Himalaya, India. *Quart. Res.*, v. 70, p. 68-80.
- Wasson, R.J., Juyal, N., Jaiswal, M., McCulloch, M., Sarin, M.M., Jain, V., Srivastava, P. and Singhvi, A.K., 2008. The Mountain-Lowland Debate: deforestation and Sediment Transport in the Upper Ganga Catchment. *Environ. Manag.*, v. 88, p. 53-61.
- Yogesh, P., Ganesh Prasad., Choubey, V.M. and Ramola, R.C., 2009. Geohydrological control on radon availability in groundwater. *Radiation Measurements*, v. 44, p. 122-126.
- Yogesh, P., Ganesh, P., Gusai, G.S., Ramola, R.C. and Choubey, V.M., 2008. Soil exhalation rate from soil samples of south Kumaun, Lesser Himalaya, India, *Radiation Measurements*, v. 43, p. S369-S374.
- 2009-10**
- Bhandari, A., Mohabey, D.M., Bajpai, S., Tiwari, B.N. and Pickford, M., 2009. Early Miocene mammals from central Kutch (Gujarat), Western India: implications for geochronology, biogeography, eustacy and intercontinental dispersals, *N. Jb. Geol. Paläont. Abh.*, v. 256, p. 69-97.
- Chaujar, R.K., 2009. Climate change and its impact on the Himalayan glaciers – a case study on the Chorabari glacier, Garhwal Himalaya, India, *Current Science*, v. 96, p. 703-708.
- Choubey, V. M., Kumar, Naresh. and Arora, B.R., 2009. Precursory signatures in the radon and geo-hydrological borehole data for M4.9 Kharshali Earthquake of Garhwal Himalaya. *Sci.of Total Environ.*, v. 407, p. 5877-5883.
- Ghosh, S. K., Ray, Y. and Sinha S. 2010. Soft sediment deformation structures: Seismites or penecontemporaneous, a case study from the Proterozoic Lesser Himalaya, India, *Current Science*, v. 98(2), p. 247-253.
- Gupta, V., 2009. Non Destructive Testing of some Higher Himalayan rocks. *Bull Eng. Geol. Env.*, v. 68, p. 409-416.
- Gupta, V. Sharma, R. and Sah, M. P., 2009. An evaluation of surface hardness of natural and modified rocks using Schmidt hammer: Study from northwestern Himalaya. *Geografiska Annaler*, v. 91A, p. 179-188.
- Hazarika, D. Baruah, S. and Gogoi, N. K., 2009. Attenuation of coda waves in North Eastern Region of India. *Jour. Seism.*, v. 13, p. 141-160.
- Jayangondaperumal, R., Dubey, A.K. and Sen, K., 2010. Mesoscopic and magnetic fabrics in arcuate igneous bodies: an example from the Mandi-Karsog pluton, Himachal Lesser Himalaya, *Geol. Mag.*, Cambridge University Press, p. 1-13.
- Jayangondaperumal, R., Dubey, A.K., Senthil Kumar, B., Wesnously, and Sangode, S. J., 2009. Magnetic fabrics indicating Late Quaternary seismicity in the Himalayan foothills, *Internat. Jour. Earth Sci.*, v. 99(1), p. 265-278.
- Kohn, M. J., Paul, S.K. and Corrie, S.L., 2010. The lower Lesser Himalayan Sequence: a Paleoproterozoic continental arc margin to the northern Indian Plate, *Jour. Geol. Soc. Amer. Bull.*, v. 122 (3/4), p. 323-335.
- Kumaravel, V., Sangode, S. J., Siddaiah, N. S. and Kumar, R., 2010. Magnetic susceptibility relationship with Munsell Soil color and geochemical parameters in the paleosol sequences from the Himalaya: An example from Pliocene-Pleistocene Siwalik paleosols NW Himalaya, India, *Geoderma*, v. 154, p. 267-280.
- Mahajan, A.K., 2009. NEHRP soil classification and estimation of 1-D site effect of Dehradun Fan deposits using Shear Wave Velocity. *Eng. Geol.*, v. 104, p. 232-240.
- Mahajan, A.K., Thakur, V.C., Sharma, M.L. and Chauhan, M., 2009. Probabilistic Seismic Hazard Map of NW Himalaya and its Adjoining Area, India. *Natural Hazard*, v. 53, p. 443-457.
- Mathur, N.S., Juyal, K.P. and Kumar, K., 2009. Larger foraminiferal biostratigraphy of lower Paleogene successions of Zanskar Tethyan and Indus-Tsangpo Suture zones, Ladakh, India in the light of additional data. *Him. Geol.*, v. 30, p. 45-68.
- Misra, D. K. and Srivastava, P., 2009. River response to continuing movements along the active faults in the Siang Valley, North-Eastern Himalaya, India, *Zietschrift Fur Geomorphologie*, v. 53,

- p. 455-468.
- Mollai, H. Sharma, R. and Georgia, Pe-Piper, 2009. Copper mineralization around the Ahar Batholith, north of Ahar (NW Iran): evidence for fluid evolution and the origin of the skarn ore deposit, *Ore Geol. Rev.*, v. 35, p. 401-414.
- Mukherjee, B.K. and Sachan, H.K., 2009. Fluids in coesite-bearing rocks of the Tso Morari Complex, NW Himalaya: evidence for entrapment during peak metamorphism and subsequent uplift, *Geol. Mag.*, v. 146 (6), p. 876-889.
- Myrow, Paul M., Hughes, Nigel C., Searle, Michael P., Fanning, C.M., Peng, S.C. and Parcha S.K., 2009. Stratigraphic correlation of Cambrian-Ordovician deposits along the Himalaya: Implications for the age and nature of rocks in the Mount Everest region. *Geol. Soc. Amer. Bull.*, v. 121, p. 323-332.
- Pandey, P., Kumar, R., Suresh, N., Sangode, S.J. and Pandey, A.K., 2009. Soft sediment deformation in contemporary reservoir sediment: a repository to recent major earthquake events in Garhwal Himal. *Jour. Geol.*, v. 117, p. 200-209.
- Peng, S.C., Hughes, N.C., Noel, A. H., Bryan, K. S., Xuejian Zhu, Myrow, P. M. and Parcha, Suraj, K., 2009. Cambrian Trilobites from the Parahio and Zanskar valleys, Indian Himalaya. *Jour. Palaeont.*, v. 83(6), p. 1-95.
- Phartiyal, B., Sharma, A., Srivastava, P. and Ray, Y., 2009. Chronology of relict lake deposits in the Spiti River, NW Trans Himalaya: Implications to Late Pleistocene-Holocene climate-tectonic perturbations. *Geomorphology*, v. 108, p. 268-272.
- Pickford, M., Bhandari, A., Tiwari, B.N. & Mohabey, D.M., 2009. Miocene terrestrial mammals from circum-Indian Ocean, focus on eustacy and Himalayan orogenesis. *Him. Geol.*, v. 30, p. 193-198.
- Prasad, Yogesh, Prasad, Ganesh, Gusain, G.S., Choubey, V.M. and Ramola, R. C., 2009. Seasonal variation of radon emission from soil and water. *Ind. Jour. Phys.*, v. 83, p.1001-1010.
- Rajesh, S. and Majumbar, T.J., 2010. Geoid height versus topography of the Northern Ninetyeast Ridge: Implications on crustal compensation, *Marine Geophys. Res.*, DOI 10.1007/s 11001-010-9088-7.
- Rameshwar, Rao, D. and Rai, Hakim, 2009. Geochemical studies of granitoids from Shyok tectonic zone of Khardung-Panamik section, Ladakh, India. *Jour. Geol. Soc. India*, v. 74, p.363-374.
- Rameshwar, Rao, D. and Sharma, Rajesh., 2009. Petrogenesis of the granitoid rocks from Askot crystallines, Kumaun Himalaya. *Jour. Geol. Soc India*, v. 74, p. 363-374.
- Rose, K.D., Rana, R.S., Sahni, A., Kumar, K., Missiaen, P., Singh, L. and Smith, T., 2009. Early Eocene Primates from Gujarat, India. *Jour. Human Evolution*, v. 56, p. 366-404.
- Rose, K.D., Rana, R.S., Sahni, A., Kumar, K., Singh, L. and Smith, T., 2009. First tillodont mammal from India: additional evidence for an early Eocene faunal connection between Europe and India? *Acta Palaeont. Polonica*, v. 54, p. 351-355.
- Saraf, A. K., Rawat, Vineeta, Choudhury, Swapnamita, Dasgupta, Sudipta and Das, J. D., 2009. Advances in understanding of the mechanism for generation of earthquake thermal precursors detected by satellites, *Internat. Jour. Appl. Earth Observations and Geoinformation*, v. 11(6), p. 373-379
- Siddaiah, N.S. and Kumar, K., 2009. Discovery of Minamiite from the Deccan volcanic province, India and implications for Martian surface exploration. *Curr. Sci.*, v. 97(11), p. 1664-1669.
- Singh, A.K. and Vallinayagam, G., 2009. Radioactive element distribution and rare-metal mineralization in Anorogenic acid volcano-plutonic rocks of the Neoproterozoic Malani Felsic Province (MFP), Western Peninsular India. *Jour. Geol. Soc. India*, v. 73, p. 837-853.
- Suresh, N. and Kumar, R., 2009. Variable period of aggradation and termination history of two late Quaternary alluvial fans in the Soan dun, NW sub Himalaya: impact of tectonic and climate. *Him Geol.*, v. 30, p. 155-165.
- Taverne, L., Kumar, K. and Rana, R.S., 2009. Complement to the study of the Indian Paleocene osteoglossid genus Taverneichthys (Teleostei, Osteoglossomorpha). *Bull. de l'Institut Royal des Sci. Naturelles de Belgique, Sci. de la Terre*, v. 79, p. 155-160.
- Tiwari, M. and Pant, I., 2009. Microfossils from the Neoproterozoic Gangolihat Formation, Kumaun Lesser Himalaya: Their stratigraphic and evolutionary significance, *Jour. Asian Earth Sci.*, v. 35, p. 137-149.
- Tewari, V.C., 2009. World Summit on Ancient Microscopic Fossils, Episodes, v. 32, p. 212-213.
- Whiso, K., Tiwari, B.N., Bajpai, S., Cooper, L.N. and Thewissen, J.G.M., 2009. A fossil mammal from marine Eocene strata (Jaintia Group) of the Mikir hills, Assam, northeastern India, *Jour. Palaeont. Soc. India*, v. 54(1), p. 111-114.

## 2010-11

- Bera, M.K., Bhattacharya, K., Sarkar, A., Samanta, A., Kumar, K. and Sahni, A., 2010. Oxygen isotope analysis of bone and tooth enamel phosphate from Paleogene sediments: experimental techniques and initial results. *Jour. Geol. Soc. India*, v. 76, p. 275-282.
- Bhambri, R., Bolch, T. and Chaujar, R.K., 2011. Mapping of debris-covered glaciers in the Garhwal Himalayas using ASTER DEMs and thermal data. *Internat. Jour. Remote Sensing*, v. 32, p. 8095-8119.
- Bhaumik A.K., Gupta, A.K., and Thomas, E., 2011. Blake Outer Ridge: Late Neogene variability in paleoceanography and deep-sea biota. *Palaeogeogr. Palaeoclim. Palaeoecol.*, v. 302, p. 435-451.
- Chaujar, R.K., 2010. Lichenometric studies and Pindari Glacier Retreat, *Current Science*, v. 99(10), p. 1306-1307.
- Choubey, V.M., Ahmad, Irfan, Kamra, Leena and Ramola, R.C., 2010. Radon variation in soil and groundwater of Bhilangana Valley, Garhwal Himalaya, India. *Japanese Jour. Health Phys.*, v. 45(3), p. 278-283.
- Choubey, V.M., Arora, B. R., Barbosa, S. M., Kumar, Naresh and Kamra, L., 2011. Seasonal and daily variation of Radon at 10 m depth in borehole, Lesser Garhwal Himalaya, India. *App. Radiation and Isotopes*, v. 69, p. 1070-1078.
- Cooper, L.N., Thewissen, J.G.M., Bajpai, S. and Tiwari, B.N., 2011. Postcranial morphology and locomotion of the Eocene raoellid *Indohyus* (Artiodactyla: Mammalia). *Historical Biology*, v. 24, p. 279-310.
- Dubey, A. K., 2010. Role of inversion tectonics in structural development of the Himalaya, *Jour. Asian Earth Sci.*, v. 39, p. 627-634.
- Dudkin, F., Rawat, Gautam, Arora, B.R., Koreponov, V., Leonteyav, O.

- and Sharma, A.K. 2010. Application of Polarisation ellipse technique for analysis of ULF magnetic fields from two distant stations in Koyana-Warna seismoactive region, western India. *Nat. Hazards Earth Syst. Sci.*, v. 10, p.1513-1522.
- Dutta, S., Mallick, M., Kumar, K., Poinar, G., Mann, U. and Greenwood, P.F., 2011. Terpenoid composition and botanical affinity of Cretaceous resins from India and Myanmar. *Intern. Jour. Coal Geology*, v. 85, p. 49-55.
- Guhey, R., Sinha, D. and Tewari, V.C., 2011. Meso- Neoproterozoic stromatolites from the Indravati and Chhattisgarh basins, Central India. *In: Stromatolites*, (V.C. Tewari and J. Seckbach, Eds.), Interaction of Microbes with Sediments, Cellular Origin. Life in Extreme Habitats and Astrobiology, Springer Science - Business Media B.V., v. 18, p. 21-42.
- Gupta, A. K., Mohan, K. Sarkar, S., Clemens, S.C., Ravindra, R. and Uttam, R.K., 2011. East–West similarities and differences in the surface and deep northern Arabian Sea records during the past 21 Kyr, *Palaeogeogr. Palaeoclim. Palaeoecol.*, v. 301, p. 75-85.
- Gupta, Vikram and Mazari, R.K., 2010. Slope Stability study of the proposed Madhyamaheshwar Small Hydropower Project (MSHP), District Rudrapur, Uttarakhand, India. *Landslides*, v. 3, p. 17-24.
- Gupta, Vikram, Sharma, Ruchika and Sah, M.P., 2011. Surface weathering of gneiss, northwestern higher Himalaya, India, *Quart. Jour. Eng. Geol. and Hydrogeol.*, v. 44, p. 135-140.
- Kayal, J.R., Arifiev, S.S., Baruah, S., Tatarovsian, R., Gogoi, N., Sanjoun, M., Gautom, J.L., Hazarika, Devajit and Borah, D., 2010. The 2009 Bhutan and Assam felt earthquakes (Mw 6.3 and 5.1) at the Kopili fault in the northeast Himalaya region, *Geomatics. Nat. Hazards and Risk*, v. 1, p. 273-281.
- Kothiyari, G.C., Pant, P.D., Joshi, M., Luirei, Khayingshing and Malik, J.N., 2010. Active faulting and deformation of Quaternary landform Sub-Himalaya, India. *Geochronometria*, v. 37, p. 63-71.
- Krishnamurthi R., Sen, A. K., Kumar, Pradeep., T. and Sharma, Rajesh., 2010. Genesis of gold mineralization in the southern granulite terrain of Peninsular India. *In: Gold Metallogeny in India and Beyond* (Mihir Deb and Richard J. Goldfarb, Eds), Narosa Publications, p. 222-233.
- Kumar, K., Rose, K.D., Rana, R.S., Singh, L., Thierry, S. and Sahni, A., 2010. Early Eocene artiodactyls (Mammalia) from western India. *Jour. Verteb. Palaeont.*, v. 30(4), p. 1245-1274.
- Kumar, K., Singh, H. and Rana, R.S., 2011. Ichnospecies *Teredolites longissimus* and teredinid body fossils from the Early Eocene of India—taphonomic and palaeoenvironmental implications. *Ichnos*, v. 18(2), p. 57-71.
- Lokho, K., Raju, D.S.N. and Azmi, R.J., 2011. Palaeoenvironmental and Biostratigraphic Significance of Uvigerinids and other Foraminifera from the Bhuban Formation, Assam-Arakan Basin, Mizoram. *Jour. Geol. Soc. India*, v. 77, p. 252-260.
- Lyubushin, A.A., Arora, B.R. and Kumar, Naresh, 2010. Investigation of seismicity in western Himalaya. *Russ. J. Geophy. Research*, v. 11(1), p. 27-34.
- Mahajan, A.K., Galiana-Merino, J.J., Lindholm, C., Arora, B.R., Mundepi, A. K., Rai, N. and Chauhan, Neetu, 2011. Characterization of the Sedimentary Cover at the Himalayan Foothills using Active and Passive Seismic Techniques. *Jour. Applied Geophy.*, v. 73, p. 196-206
- Mahajan, A.K., Thakur V.C., Lal, Mukut and Chauhan, Mukesh, 2010. Probabilistic Seismic Hazard Map of NW Himalaya and its adjoining area, India. *Natural Hazard*, v. 53, p. 443-457.
- Mayr, G., Rana, R.S., Rose, K.D., Sahni, A., Kumar, K., Singh, L. and Smith, T., 2010. Quercypitta-like birds from the early Eocene of India (Aves, ?Psittaciformes). *Jour. Verteb. Palaeont.*, v. 30, p. 467-478.
- Mega Daga, M., Rameshwar Rao, D. and Rai, Hakim, 2010. Granite magmatism and crustal evolution of the Shyok-Darbuk corridor of NE Ladakh, India. *IUGS Episodes*, v. 33(3), p. 169-185.
- Meena, N. K., Maiti, S., and Srivastava, A., 2011. Discrimination between anthropogenic (pollution) and lithogenic magnetic fraction in urban soils (Delhi, India) using environmental magnetism. *Jour. Appl. Geophys.*, v. 73, p. 21-129.
- Mrinalinee Devi, R. K., Bhakuni S. S., and Bora, P. K., 2010. Neotectonic study along mountain front of northeast Himalaya, Arunachal Pradesh, India, *Environ Earth Sci*, v. 63(4), p. 751-762.
- Mrinalinee Devi, R.K., Bhakuni, S.S. and Bora, P.K., 2011. Tectonic implication of drainage set-up in the Sub-Himalaya: a case study of Papumpare district, Arunachal Himalaya, India. *Geomorphology*, v. 127, p. 14-31.
- Mukhopadhyay, S., Sharma, J., Del-Pezzo, E. and Kumar, Naresh, 2010. Study of attenuation mechanism for Garhwal-Kumaon Himalayas from analysis of coda of local earthquakes. *Physic. Earth. Planet. Internat.*, v. 180, p. 7-15.
- Myrow, Paul M., Hughes, Nigel C., Goodge, John W., Fanning, C. Mark., Williams, Ian S., Peng, Shanchi, Bhargava, Om N., Parcha, Suraj K., and Pogue, Kevin R., 2010. Extraordinary transport and mixing of sediment across Himalayan central Gondwana during the Cambrian–Ordovician. *Geol. Soc. Amer. Bull.*, v. 122(9/10), p. 1660-1670.
- Nandini P. and S. S. Thakur, 2011. Metamorphic Evolution of Lesser Himalayan Crystalline Sequence (LHCS), Siyom Valley, NE Himalaya, India. *Jour. Asian Earth Sci.*, v. 40, p. 1089-1100.
- Parcha, S.K. and Pandey, S. 2011. Devonian Ichnofossils from the Farakah Muth Section of the Pin Valley, Spiti Himalaya. *Jour. Geol. Soc. India*, v. 78, p. 263-270.
- Parcha, S.K. and Pandey, S., 2011. Ichnofossils and their significance in the Cambrian successions of the Parahio Valley in the Spiti Basin, Tethys Himalaya, India, *Jour. Asian Earth Sci.*, v. 42, p. 1097-1116.
- Philip, G., Suresh, N., Bhakuni, S.S. and Gupta, V., 2011. Paleoseismic investigation along Nalagarh Thrust: Evidence of late Pleistocene earthquakes in Pinjaur Dun, Northwestern Sub Himalaya, India, *Jour. Asian Earth Sci.*, v. 40, p.1056-1067.
- Prakash, D., Prakash, S. and Sachan, H.K., 2010. Petrological evolution of the high pressure and ultrahightemperature mafic granulites from Karur, southern India: evidence for decompressive and cooling retrograde trajectories. *Mineral. Petrol.*, v. 100 (1-2), p. 35-53.
- Ramola, R.C., Prasad, G., Gusai, G.S., Rautala, B.S., Choubey, V.M., Vidyasagar, D., Tokonami, S., Sorimachi, A., Sahoo, S.K., Janik, M. and Ishikawa, T., 2010. Preliminary indoor thoron measurements in high radiation background area of southeastern

- coastal Orissa, India. *Radiation Proct. Dosimetry*, v. 141(4), p. 379-382.
- Rai, Santosh K., Singh, S. K. and Krishnaswami, S., 2010. Chemical weathering in the Plain and Peninsular sub-basins of the Ganga: Impact on Major ion chemistry and Elemental Fluxes. *Geochim. Cosmoch. Acta*, v. 74(8), p. 2340-2355.
- Rajesh, S. and Majumdar, T. J., 2010. Geoid versus Topography of the Northern Ninetyeast Ridge: Implications on crustal compensation. *Marine Geophys. Res.*, v. 30(4), p. 251-264.
- Ray, Yogesh and Srivastava, Pradeep, 2010. Widespread aggradation in the mountainous catchment of the Alaknanda-Ganga River System: Timescales and implications to Hinterland-foreland relationships. *Quarter. Sci. Rev.*, v. 29, p. 2238-2260.
- Sachan, H.K., Kohn, M. J., Saxena, A. and Corrie, S.L., 2010. The Malari leucogranite, Garhwal Himalaya, northern India: chemistry, age, and tectonic implications. *Geol. Soc. Amer. Bull.*, v. 122(11/12), p. 1865-1876.
- Sanyal, P, Sarkar, A., Bhattacharya, S.K., Kumar, R., Ghosh, S.K. and Agarwal, S., 2010. Intensification of monsoon, microclimate and asynchronous C4 appearance: isotopic evidence from the Indian Siwalik sediments. *Palaeogeogr. Palaeoclim. Palaeoecol.*, v. 296, p. 165-173.
- Sharma, R., Joshi, P. and Verma, P., 2010. Genetic issues of some of the non-metallic minerals in Lesser Himalaya. *Iranian Jour. Earth Sci.*, v. 2, p. 168-172.
- Sharma, Ruchika, Gupta, V., Arora, B. R. and Sen, K., 2010. Petrophysical properties of the Himalayan Granitoids: Implication on composition and source. *Tectonophysics*, v. 497, p. 23-33.
- Shashidhar, D., Kumar, Naresh, Mallika, K. and Gupta, H. K., 2010. Characteristics of Seismicity Pattern prior to the M ~ 5 Earthquakes in the Koyna Region, Western India – Application of the RTL Algorithm. *Episodes*, v. 33(2), p. 83-90.
- Sinha, S., Suresh, N., Kumar, R., Dutta, S. and Arora, B. R., 2010. Sedimentologic and geomorphic studies on the Quaternary alluvial fan and terrace deposits along the Ganga exit. *Quarter. Internat.*, v. 227, p.87-103.
- Singh, H., Prasad, M., Kumar, K. and Singh, S., 2011. Paleobotanical remains from the Paleocene–Lower Eocene Vagadkhol Formation, western India and their climatic and phytogeographic implications. *Palaeoworld*, v. 20, p. 332-356.
- Singh, H., Prasad, M., Kumar, K., Rana, R.S. and Singh, S.K., 2010. Fossil fruits from Early Eocene Vastan Lignite, Gujarat, India: taphonomic and phytogeographic implications. *Current Science*, v. 98 (12), p. 1625-1632.
- Singh, Keser, 2010. Tectonic evolution of Kishtwar Window with respect to the Main Central Thrust, northwest Himalaya, India. *Jour. Asian Earth Sci.*, v. 39, p. 125-135.
- Thakur, V.C., Jayangondaperumal, R. and Malik, M.A., 2010. Redefining Medicott-Wadia main boundary fault from Jhelum to Yamuna: an active fault strand of the main boundary thrust in northwest Himalaya. *Tectonophysics*, v. 489, p. 29-42.